

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS	PA Required	Criteria
----------	--------------------	------------	-----------------	--------------------	------------	---------------------	-------------	----------

PDL Effective November 8, 2024

***PLEASE NOTE: For a search box hit Ctrl F**

*** PLEASE NOTE: All cost effective generics applicable to DEL are considered PREFERRED Drugs. "BASIC" Covered Drugs are bolded with the Coverage Indicator of "MC / DEL".**

General Criteria for all PDL categories- For more information or help using the PDL, providers may call 1-888-445-0497; members should call 1-866-796-2463. To access PDL and PA materials via the internet: www.mainearepdl.org

A: Preferred Drugs- Unless otherwise specified, preferred drugs are available without prior authorization. Step order may apply for preferred drugs in some drug categories as indicated on the PDL. (See item "D" below for explanation of step order.)

C: Adequate Drug Trials- 1. The minimum trial period for each preferred and step order drug is two weeks, unless otherwise stated within specific PDL drug categories; trials with less than a two week duration will be reviewed on a case-by-case basis; 2. A trial will not be considered valid if preferred or non-preferred products were readily available (by override, individual purchase, samples, etc.); 3. Certain drug trials, such as with controlled substances, may require evidence that the preferred drugs were actually tried (example: with random pill counts and with random urine drug tests, using the methods of GC/MS with no lower threshold); 4. Adequate trials require documentation of attempts to titrate dose of preferred agents toward desired clinical response. 5. Adequate trials include prevention/treatment of common adverse effects associated with preferred agents (example: antinausea, antipruritics, etc.)

D: Step Order- When numbers appear in the "step order" column, it means drugs in this category must be used in the order specified, with the lower numbers having preference over the higher numbers. Chart notes should be provided to confirm drug trials that do not appear in the member's MaineCare drug profile.

E. The Department will institute strategies to ensure cost effectiveness through the use of an enhanced Drug Benefit Preferred brand drugs will no longer be preferred in any PDL drug category where preferred generic drugs are also available. It is expected that preferred generics will be used prior to any preferred brands. This will be operated as a form of step care. Preferred brands in these categories will require prior authorization for these high utilization / high cost members.

F: Brand Name Medication Requests- (Must be submitted on the Brand Name PA request form)- According to MaineCare Benefits Manual Chapter II (80.07-5), when medically necessary covered brand-name drugs have an A-rated generic equivalent available, the most cost effective medically necessary version will be approved and reimbursed, since the brand-name and A-rated generic drugs have been determined by the FDA to be chemically and therapeutically equivalent. The Bureau does not make determinations as to whether or not a generic drug is clinically inferior or inequivalent to its brand version. This is the proper role of the FDA. Physicians should submit their reports of generic inequivalence directly to the FDA via the MEDWATCH.

G: PA requests for non- FDA Approved Indications- Decisions will be made on a case-by-case basis until the DUR committee is able to review the evidence and make a recommendation. Interim approvals and DUR recommendations for approval of a drug for a non- FDA approved indication will require a minimum of two published, peer reviewed, non contradicted, double- blind, placebo-controlled randomized clinical studies establishing both safety and efficacy.

H: Dose Consolidation Requirements- Some drugs may also be affected by dose consolidation requirements. Please see Dose Consolidation List and/or Splitting Tables provided in the PDL.

I. Trials from Multiple Drug Classes - Trial/failure/intolerance to preferred agents from multiple classes within the same category or other categories of drugs may be required prior to the approval of non-preferred agents (e.g., Cymbalta, Zofran, Elidel and others).

J. Drug-specific PA Forms- Drug-specific PA forms contain medical necessity documentation requirements and/or criteria that may not be repeated in the PDL. Drug-specific PA forms may be obtained on the web at www.mainearepdl.org.

K. PA Exemptions for Prescribers- According to MaineCare Benefits Manual Chapter II (80.07-4), providers may receive a three (3) month exemption from prior authorization requirement for certain categories of drugs when they demonstrate high compliance with the Department's PDL. The Department will notify providers in writing which drug categories are included and what dates apply to the exemption. If a provider loses his/ her exemption, members who previously were not required to obtain a PA while the prescriber was exempt will be required to do so, and criteria for approval of that medication will need to be met.

L: Drug-Drug Interactions (DDI)- The DUR Committee has implemented new drug-drug interaction edits requiring prior authorization. Several drug-drug combinations and PDL drug categories are affected by new PA requirements. These will be indicated in the PDL with DDI notation. Please see the DDI document provided in the PDL.

ASSORTED ANTIBIOTICS

BETA-LACTAMS / CLAVULANATE COMBO'S	MC/DEL	AMOXICILLIN	MC/DEL	AUGMENTIN ³	3. Chewable 125mg & 250mg and Solution 125mg/5ml and 250mg/5ml available without PA. 4. Use preferred generic amoxicillin/clavulanate potassium alternatives. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Ampicillin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI.
	MC/DEL	AMOXICILLIN/POTASSIUM CLA CHEW	MC/DEL	AUGMENTIN XR TB12 ⁴		
	MC/DEL	AMOXICILLIN/POTASSIUM CLA SUSR				
	MC/DEL	AMOXICILLIN/POTASSIUM CLA TABS				
	MC/DEL	AMPICILLIN				
	MC	BICILLIN L-A SUSP				
	MC/DEL	DICLOXACILLIN SODIUM CAPS				
	MC	OXACILLIN SODIUM SOLR				
	MC/DEL	PENICILLIN V POTASSIUM				
	MC	TIMENTIN SOLR				
MC	UNASYN SOLR					
MC/DEL	ZOSYN					
CEPHALOSPORINS	MC/DEL	CEFADROXIL HEMIHYDRATE	MC	CEDAX	1. Both brand and generic are clinically non-preferred. 2. Dosing limits apply, please see Dosage Consolidation List. 3. Approvals will only be considered for patients 18 years of age or older who have limited or no alternative	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Vantin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non
	MC/DEL	CEFAZOLIN SODIUM SOLR	MC/DEL	CEFACTOR ¹		
	MC/DEL	CEFDINIR	MC/DEL	CEFADROXIL MONOHYDRATE TABS		
	MC/DEL	CEFEPIME	MC/DEL	CEFIXIME SUS		
	MC/DEL	CEFPODOXIME	MC/DEL	CEPHALEXIN TABS		
	MC/DEL	CEFPODOXIME PROXETIL SUS	MC	CEPHALEXIN 750MG CAPS		
	MC/DEL	CEFPODOXIME PROXETIL TAB	MC/DEL	CEFTIN		
	MC/DEL	CEFIXIME 400MG ² CAP	MC	DAXBIA		
	MC/DEL	CEFPROZIL	MC	FETROJA ³		

							Arikayce will require clinical PA to confirm MAC lung disease and for use in adults who have limited or no alternative treatment options. Zemdri will be reserved for patients with limited or no alternative treatment of care.
ANTI-MYCOBACTERIALS / ANTI-TUBERCULOSIS	MC/DEL MC/DEL MC/DEL MC/DEL		ETHAMBUTOL HCL TABS MYAMBUTOL TABS RIFABUTIN CAPS RIFAMPIN	MC/DEL MC/DEL MC		MYCOBUTIN CAPS PRETOMANID RIFADIN CAPS	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Pretomanid is indicated as part of a combination regimen with bedaquiline and linezolid for the treatment of adults with pulmonary extensively drug resistant (XDR) or treatment-intolerant or non-responsive multidrug-resistant (MDR) tuberculosis (TB). Approval of this indication is based in limited clinical safety and efficacy data. This drug is indicated for use in a limited and specific population of patients. DDI: Preferred rifampin will be non-preferred and require prior authorization if it is currently being used in combination with either Pradaxa or Latuda.
ANTIMALARIAL AGENTS	MC/DEL MC MC/DEL MC/DEL		DARAPRIM TABS KRINTAFEL ² MEFLOQUINE HCL TABS QUININE SULFATE	MC MC/DEL MC/DEL MC MC MC/DEL		ARALEN TABS CHLOROQUINE PHOSPHATE TABS ³ HYDROXYCHLOROQUINE TABS ³ ISONARIF ¹ MALARONE TABS PLAQUENIL TABS	Use PA Form# 20420 1. Ingredients available as preferred without PA. 2. Krintafel is preferred for ≥ 16 years of age. 3. Established users will be grandfathered Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid coadministration of Krintafel® with Organic Cation Transporter 2 (OCT2) and Multidrug and Toxin Extrusion (MATE) substrates (e.g. dofetilide, metformin).
ANTHELMINTICS	MC/DEL MC/DEL MC/DEL		ALBENDAZOLE PRAZIQUANTEL TAB STROMECTOL TABS	MC MC MC/DEL		ALBENZA TABS EMVERM BILTRICIDE TABS	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIBIOTICS - MISC.	MC MC MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC		AZACTAM SOLR COLY-MYCIN-M SOLR COLISTIMETHATE SODIUM SOLR FIRVANQ ⁴ FUROXONE TABS METRONIDAZOLE ¹ PENTAMIDINE ISETHIONATE SOLR SOLOSEC TRIMETHOPRIM TABS VANCOMYCIN 5GM INJ. VANCOMYCIN CAPS XIFAXAN 200mg	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC MC		AEMCOLO COLISTIMETHATE SODIUM SOLR CAYSTON ³ FLAGYL CAPS FLAGYL TABS FLAGYL ER TBCR KETEK LIKMEZ METRONIDAZOLE 375MG CAPS ¹ METRONIDAZOLE 750MG TABS ¹ NEBUPENT SOLR REBYOTA ⁵ TINDAMAX VANCOMYCIN 10GM INJ. ² XENLETA XIFAXAN VOWST ⁵	1. 375mg caps and 750mg tabs are non-preferred. Please use available preferred strengths(250mg & 500mg tabs) to obtain required dose without PA. 2. Please use multiple 5gm which are preferred to obtain dose without PA. 3. Clinical PA is required to establish CF diagnosis and medical necessity. Prior trial and failure of preferred Tobi before approval will be granted. 4. Quantity limit of one per 150ml bottle. 5. For the treatment of patients 18 years of age and older. Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. 1. For macrolide resistant infections when quinolones inappropriate DDI: Ketek is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either Enblex 15mg or Vesicare 10mg or carbamazepine. Cayston is only indicated to improve respiratory symptoms in CF patients with Pseudomonas aeruginosa. Dosing limits, as should be given TID X28 days (followed by 28 days OFF Cayston therapy). A bronshodilator should be used before administration of Cayston. Xenleta will be considered for the treatment of adults with community-acquired bacterial pneumonia (CABP) caused by the following susceptible microorganisms: Streptococcus pneumoniae, Staphylococcus aureus (methicillin-susceptible isolates), Hemophilus influenzae, Legionella pneumophila, Mycoplasma pneumoniae, and Chlamydia pneumoniae. Vowst: To prevent the recurrence of Clostridioides difficile infection (CDI) in individuals 18 years of age and older following antibacterial treatment for recurrent CDI (rCDI). Likmez: patient has a medical necessity for a non-solid oral dosage form. Rebyota: For the prevention of recurrence of Clostridioides difficile infection (CDI) in individuals 18 years of age and older following antibiotic treatment for recurrent CDI. The limitation of use is that Rebyota® is not indicated for treatment of CDI.
CARBAPENEMS				MC MC MC/DEL MC/DEL		INVANZ SOLR MERREM SOLR PRIMAXIN RECARBRIO	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

LINCOSAMIDES / OXAZOLIDINONES / LEPROSTATICS	MC/DEL MC/DEL MC/DEL MC MC/DEL		CLEOCIN SOLN CLEOCIN SUSR CLINDAMYCIN HCL 150CAPS DAPSONE TABS LINEZOLID 600mg TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL	8 8 8 8 8 9 9	CLEOCIN CAPS CLINDAMYCIN HCL 300CAPS ¹ SIVEXTRO VIBATIV LINEZOLID TABS ZYVOX SUSR ZYVOX TABS	1. Use multiple 150's for Clindamycin instead of 300's. 2. Quantity limit of 14 days supply within a 60day period. Use PA Form# 30820 for Zyvox & Vibativ Use PA Form# 20420 for all others	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. For Zyvox or Vibativ, please see the criteria listed in the Antibacterial Antibiotics PA form.	
ANTI INFECTIVE COMBO'S - MISC.	MC/DEL MC/DEL MC/DEL MC/DEL		ERYTHROMYCIN/SULF SUSR SEPTRA/DS TABS SULFAMETHOXAZOLE/TRIMETH TRIMETHOPRIM/SULFAMETHOXA	MC MC		BACTRIM DS TABS VABOMERE ¹	Use PA Form# 20420 1. For the treatment of patients ≥ 18 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
ANTIPROTOZOALS	MC/DEL MC/DEL		BENZNIDAZOLE ² LAMPIT ²	MC		ALINIA ¹	1. Alina is preferred for children less than 12 years of age. 2. Clinical PA required for appropriate diagnosis. Use PA Form# 20420	Benznidazole is indicated for pediatric patients 2 to 12 years of age for the treatment of Chagas disease (American trypanosomiasis) caused by Trypanosoma cruzi.	
ANTI - FUNGALS									
ANTIFUNGALS - ASSORTED	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ANCOBON CAPS FLUCONAZOLE ¹ KETOCONAZOLE TABS ⁷ NYSTATIN TERBINAFINE TABS ⁴ VORICONAZOLE TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	6 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	LAMISIL TABS ⁴ ITRACONAZOLE BREXAFEMME CRESEMBA ⁹ GRIFULVIN V TABS GRISEOFULVIN SUSP GRISEOFULVIN ULTRAMICROSI TABS GRIS-PEG TABS REZZAYO ⁹ SPORANOX SOLN ² SPORANOX PULSEPAK CAPS ³ SPORANOX CAPS ³ DIFLUCAN ERAXIS INJ ⁶ GRIFULVIN SUSP ONMEL NOXAFIL ⁵ TOLSURA VFEND TABS VIVJOA	See quantity limit table. Non-preferred products must be used in specified step order. Continue to use Anti-Fungal PA form for non-preferred products. 1. QL--1/every 7-day period (150mg only). 2. Sporanox QL 300cc/month with PA. See quantity limit table. 3. Sporanox QL 30/month with PA. 4. Quantity limit of one tablet daily. Please see dosage consolidation list. 5. Approved if immuno suppressed/ HIV or if the member has failed a 7 day trial of a preferred antifungal therapy. 6. Eraxis will be approved if submitting with documentation that it was initiated during a hospitalization and this request is to finish the hospital course. 7. Quantity limits allowing 30 day supply without PA. PA will be required if using > 30	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The other criteria are listed on the Antifungal PA form including the required proof of a non-cosmetic fungal infection. DDI: Any Griseofulvin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI. DDI: Sporanox is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with Enablex 15mg, Vesicare 10mg, Prandin, Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI, due to a significant drug-drug interaction. DDI: Vfend is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with Warfarin. DDI: Fluconazole (except 150mg strength) will now be non-preferred and require prior authorization if it is currently being used with glimepiride (Amaryl), Enablex 15mg, or Vesicare 10mg. Diflucan is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either glimepiride (Amaryl), Enablex 15mg, or Vesicare 10mg. DDI: Fluconazole will require prior authorization if being used in combination with Plavix or Warfarin. DDI: Ketoconazole will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: Prevacid, Pantoprazole, Plavix, Onglyza, Enablex 15mg, Vesicare 10mg, Latuda, Cometriq, Tafilnar or Ormeprazole. Rezzayo: In patients 18 years of age or older who have limited or no alternative options for the treatment of candidemia and invasive candidiasis.	

ANTI - VIRALS						
ANTIRETROVIRALS	MC/DEL	ABACAVIR TABS	MC/DEL	8	ABACAVIR SOL	days.
	MC	APRETUDE	MC/DEL	8	APTIVUS	8. For children < 18, quantity limits allows 8 weeks supply without PA. PA will be required if using > than 8 weeks. If 18 and older PA will be required for any quantity. Not approving for Onychomycosis indication.
	MC/DEL	ATAZANAVIR	MC/DEL	8	CIMDUO	9. For patients ≥ 18years of age
	MC	ATRIPLA ¹	MC/DEL	8	COMBIVIR TABS	Use PA Form# 10120
	MC	BIKTARVY	MC/DEL	8	EDURANT	
	MC	CABENUVA	MC/DEL	8	EPZICOM ¹	
	MC	COMPLERA ¹	MC/DEL	8	FUZEON	
	MC/DEL	DELSTRIGO	MC/DEL	8	INTELENCE	
	MC	DESCOVY ¹	MC/DEL	8	ISENTRESS ³	
	MC	DIDANOSINE	MC/DEL	8	ISENTRESS HD	
	MC/DEL	DOVATO	MC	8	JULUCA	
	MC	EFAVIRENZ TAB	MC	8	KALETRA	
	MC/DEL	EFAVIRENZ CAP	MC/DEL	8	LAMIVUDINE SOLN	
	MC	EFAVIRENZ-EMTRICITABINE-TENOFOVIR DF TAB	MC/DEL	8	LEXIVA	
	MC	EMTRICITABINE-TENOFOVIR	MC/DEL	8	NEVIRAPINE	
	MC	EMTRIVA ¹	MC	8	NORVIR	
	MC	EPIVIR SOL	MC/DEL	8	PIFELTRO	
	MC/DEL	EVOTAZ ¹	MC	8	RETROVIR	
	MC	GENVOYA ^{1,5}	MC	8	REYATAZ	
	MC/DEL	ISENTRESS 400MG ⁶	MC/DEL	8	SELZENTRY	
	MC/DEL	ISENTRESS CHEW ³	MC	8	STAVUDINE	
	MC/DEL	ISENTRESS POWDER	MC	8	STRIBILD ¹	
	MC/DEL	LAMIVUDINE TABS	MC	8	SUNLENCA ⁵	
	MC/DEL	LAMIVUDINE/ZIDOVUDINE	MC/DEL	8	SYMFI ⁵	
	MC/DEL	LOPINAVIR-RITONAVIR SOL	MC/DEL	8	SYMFI LO ⁵	
	MC	LOPINAVIR-RITONAVIR TAB	MC/DEL	8	SYM TUZA	
	MC	ODEFSEY ¹	MC	8	TRIUMEQ ^{1,4}	
	MC/DEL	PREZCOBIX	MC/DEL	8	TRIZIVIR TABS	
	MC	PREZISTA ²	MC	8	TRUVADA ¹	
	MC/DEL	RITONAVIR TAB 100MG	MC/DEL	8	VIRACEPT TABS	
	MC	RUKOBIA ⁹	MC	8	VITEKTA	
	MC	SUSTIVA ¹	MC	8	ZERIT	
	MC	TIVICAY	MC	8	VIDEX EC	
	MC	TIVICAY PD	MC	8	VIREAD TABS ¹	
	MC	TROGARZO ⁵	MC/DEL	8	ZIAGEN TABS	
	MC	TYBOST	MC/DEL	8	ZIAGEN SOL	
	MC	VIREAD POW	MC/DEL	9	VIRAMUNE XR	
	MC/DEL	ZIDOVUDINE				
CYTO-MEGALOVIRUS AGENTS	MC	CIDOFOVIR	MC		VALCYTE TABS	Use PA Form# 20420

Fuzeon: Prescriber is either an HIV specialist provider or has consulted with one. Documentation of genotype testing is supplied and shows that there is no other potent, appropriate two or three drug oral regimen available, AND patient has a positive HIV viral load within past 6 months while on his/her current antiretroviral regimen AND the drug will be prescribed with at least two other drugs that are likely to be active based on the genotype testing.

DDI: Reyataz requires prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI .

DDI: Norvir requires prior authorization if it is currently being used in combination with either Enblex 15mg or Vesicare 10mg.

DDI: Preferred Crixivan caps requires prior authorization if it is currently being used in combination with either Enblex 15mg or Vesicare 10mg.

DDI: The concomitant use of the following drugs with **Descovy**® is not recommended: tipranavir/ritonavir, St. John's wort, and the antimycobacterials rifabutin, rifampin, or rifapentine.

DDI: Administration with the following drugs: the anticonvulsants carbamazepine, oxcarbazepine, phenobarbital, and phenytoin; the antimycobacterials rifampin and rifapentine; proton pump inhibitors such as dexlansoprazole, esomeprazole, lansoprazole, omeprazole, pantoprazole, rabeprazole; systemic dexamethasone (more than a single dose); and St. John's wort with **Odefsey** is contraindicated.

Stribild: PA required; must provider rationale as to why the member's medical need cannot be met with preferred agents, particularly Genvoya or combinations of preferred and agents AND must be antiretroviral treatment-naïve or virologically controlled on current therapy (HIV-1RNA < copies/ml) AND be HBV negative AND not be combined with other anti-retroviral agents.

DDI: Tivicay will require prior authorization is used with nevirapine, oxcarbazepine, phenytoin, phenobarbital, carbamazepine, and St. John's wort.

DDI:Aatazanavir or darunavir and the following drugs are contraindicated (due to potential for serious and/or life-threatening events or loss of therapeutic effect): alfuzosin, dronedarone, rifampin, irinotecan, dihydroergotamine, ergotamine, methylethergonovine, cisapride, St. John's wort, lovastatin, simvastatin, pimozide, nevirapine, sildenafil (when given as Revatio® for treatment of PAH), indinavir, triazolam, or PO midazolam will be non-preferred and require prior authorization if it is currently being used in combination with Tybost.

DDI: Combined P-gp, UGT1A1 and strong CYP3A inhibitors may significantly increase plasma concentrations of Sunlenca®. Concomitant administration of Sunlenca® with these inhibitors is not recommended.

Sunlenca: In combination with other antiretroviral(s) for the treatment of HIV-1 infection in heavily treatment-experienced adults with multidrug resistant HIV-1 infection failing their current antiretroviral regimen due to resistance, intolerance, or safety considerations.

	MC MC/DEL MC/DEL	FOSCARNET SODIUM GANCICLOVIR VALGANCICLOVIR	MC/DEL MC/DEL MC/DEL	FOSCAVIR LIVTENCITY ¹ PREVYMIS		1. Must show failure or contraindication to all the following ganciclovir, valganciclovir, cidofovir and foscarnet before Livtency will be approved.	on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Prevymis: Documentation that member is high-risk for CMV reactivation as defined by transplant guidelines or that there has been significant myelosuppression by one of the preferred agents. DDI: Livtency is a substrate of CYP3A4. Coadministration of Livtency® with strong inducers of CYP3A4 is not recommended, except for selected anticonvulsants.
HERPES AGENTS	MC/DEL MC/DEL	ACYCLOVIR VALACYCLOVIR HCL	MC/DEL MC MC/DEL MC MC/DEL	8 FAMCICLOVIR ¹ 8 SITAVIG 8 ZOVIRAX ¹ 8 VALTREX TABS ¹ 9 FAMVIR TABS ¹		1. Must fail Acyclovir and Valacyclovir before non-preferred products in step order. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
INFLUENZA AGENTS	MC MC MC/DEL	AMANTADINE CAPS RELENZA DISKHALER AEPB OSELTAMIVIR ¹	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL	AMANTADINE TABS FLUMADINE TABS FLUMIST RIMANTADINE HCL TABS TAMIFLU ¹ TAMIFLU SUS XOFLUZA		1. Tamiflu and Osetamivir 10 caps or 60cc's per month. Will be audited for presence of positive influenza tests in patient or family member. Use PA Form# 20420 for all others	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
IMMUNE SERUMS							
IMMUNE SERUMS	MC	HYPERRHO INJ					
HEPATITIS AGENTS							
HEPATITIS C AGENTS	MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL	SOFOSBUVIR/VELPATASVIR ² (Authorized generic labeler 72626 Asegua Therapeutics) MAVYRET ² PEGASYS KIT ¹ PEGASYS SOLN PEG-INTRON KIT ¹ RIBAVIRIN RIBASPHERE	MC/DEL MC/DEL MC MC/DEL MC MC MC MC MC/DEL	COPEGUS TABS DAKLINZA EPCLUSA ² HARVONI ² REBETOL CAPS RIBAPAK SOVALDI ² VIEKIRA PAK ² VIEKIRA XR ² VOSEVI ZEPATIER ²		1. Dosing limits apply, please see dosage consolidation list. 2. Approvals will require clinical PA. Please see the Hepatitis PA form for criteria Use PA Form #10700	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Olysio will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A4 inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin).
HEPATITIS AGENTS - MISC.			MC	ACTIMMUNE		Use PA Form# 20420	Approved for chronic granulomatous disease, osteopetrosis and idiopathic pulmonary fibrosis.
HEPATITIS B ONLY	MC/DEL MC	ENTECAVIR TENOFIVIR	MC MC MC MC	BARACLUDE HEPSERA TABS TYZEKA VEMLIDY		Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Baraclude is indicated for treatment of chronic Hep B virus (HBV) in adults with: evidence of active viral replication AND either evidence of persistent elevation in serum aminotransferases (ALT or AST) or histologically active disease, Patient is 16 years of age or older. Boxed warning: Use not recommended for those co-infected with HIV and HBV who are not also receiving highly active antiretroviral therapy (HAART). Vemlidy® remain non-preferred and require prior authorization and be available to those who have evidence of bone loss or renal insufficiency or who are unable to tolerate or who have failed on preferred medications.
RSV PROPHYLAXIS							
RSV PROPHYLAXIS			MC	SYNAGIS ¹		Use PA Form# 30120 1. MaineCare will approve Synagis PA's for start date of	Please see the criteria listed on the Synagis PA form.

November 29, 2021 for infants who meet the guidelines. PA will be approved for max of 5 doses. Maximum 1 dose/30 days. MaineCare will start accepting PAs November 1, 2021."

MS TREATMENTS

MULTIPLE SCLEROSIS - INTERFERONS	MC MC/DEL MC		AVONEX KIT ¹ BETASERON SOLR ¹ REBIF SOLN ¹	MC MC/DEL		PLEGRIDY ¹ EXTAVIA	1. Clinical PA is required to establish diagnosis and medical necessity. Use PA Form# 20430	Non-Preferred drugs must be tried in step-order and failed due to lack of efficacy or intolerable side effects before lower ranked non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
----------------------------------	--------------------	--	---	--------------	--	----------------------------------	--	---

MULTIPLE SCLEROSIS - NON-INTERFERONS	MC MC/DEL MC/DEL MC/DEL MC MC MC		COPAXONE DALFAMPRIDINE ER DIMETHYL FUMARATE CAP FINGOLIMOD CAP ² KESIMPTA ² TERIFLUNOMIDE TAB ² TYSABRI ^{1,2}	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC MC	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	AMPYRA AUBAGIO BAFIERTAM BRIUMVI GILENYA GLATOPA MAVENCLAD ³ MAYZENT OCREVUS ² PONVORY ² TASCENSO ODT ^{2,4} TECFIDERA VUMERITY ZEPOSIA	1. Providers must be enrolled in the TOUCH Prescribing program, a restricted distribution program. Clinical PA is required to establish diagnosis and medical necessity. 2. Clinical PA is required to establish diagnosis and medical necessity. 3. Due to safety profile, use of Mavenclad® is generally recommended for patients who have had an inadequate response to, or are unable to tolerate, an alternate drug indicated for the treatment of MS 4. For the treatment of patients 10 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Mavenclad will require multiple trials of preferred agents including Mayzent for secondary progressive disease. DDI: Due to significant increases in exposure to siponimod, concomitant use of Mayzent® and drugs that cause moderate CYP2C9 and moderate or strong CYP3A4 inhibition is not recommended. Ponvory: Before initiation of Ponvory® treatment, assess the following: •Complete Blood Count (CBC)- Obtain a recent (i.e. within the last 6 months) CBC, including lymphocyte count. •Cardiac Evaluation- oObtain an electrocardiogram (ECG) to determine whether pre-existing conduction abnormalities are present. In patients with certain pre-existing conditions, advice from a cardiologist should be sought and first-dose monitoring is recommended. oDetermine whether patients are taking drugs that could slow heart rate of atrioventricular (AV) conduction. •Liver Function Tests- Obtain recent (i.e. within the last 6 months) transaminase and bilirubin levels. •Ophthalmic Evaluation- Obtain an evaluation of the fundus, including the macula. •Current or prior medications with immune system effects- If patients are taking anti-neoplastic, immunosuppressive, or immune-modulating therapies, or if there is a history of prior use of these drugs, consider possible unintended additive immunosuppressive effects before starting treatment with Ponvory®. •Vaccinations- Test for antibodies to varicella zoster virus (VZV) before starting Ponvory®; VZV vaccination of antibody-negative patients is recommended prior to commencing treatment with Ponvory®. If live attenuated vaccine immunizations are required, administer at least 1 month prior to initiation of Ponvory®. •Vaccinations- Test for antibodies to varicella zoster virus (VZV) before starting Ponvory®; VZV vaccination of antibody-negative patients is recommended prior to commencing treatment with Ponvory®. If live attenuated vaccine immunizations are required, administer at least 1 month prior to initiation of Ponvory® Mayzent for Relapsing forms of MS: multiple trials of preferred agents, including an intravenous MS product. Mayzent for Active secondary progressive disease: prior trials of two preferred agents are required. Use PA Form# 20430
--------------------------------------	--	--	---	--	--	---	---	---

MULTIPLE SCLEROSIS - MISC				MC		ZINBRYTA ¹	1. The safety and efficacy of use in children under the age of 17 years have not been established. Use PA Form #20430	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists
---------------------------	--	--	--	----	--	-----------------------	--	---

ASSORTED NEUROLOGICS

NEUROLOGICS - MISC.	MC MC		BOTOX ^{2,4} DYSPORT ⁴	MC/DEL MC MC/DEL		FIRDAPSE MYOBLOC ¹ RUZURGI ³	1. Approval will be limited to Cervical dystonia.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
---------------------	----------	--	--	------------------------	--	--	---	--

				MC MC/DEL	SKYSONA ^{4,6} XEOMIN ²	<p>2. Please see botulinum PA form for additional criteria</p> <p>3. For the treatment of patients between ages 6-16 years of age.</p> <p>4. Clinical PA required.</p> <p>5. For adult patients who are anti-acetylcholine receptor (AChR) antibody positive.</p> <p>6. For the treatment of patients between ages 4-17 years of age.</p> <p>Use PA Form# 10210</p>	<p>Failed/did not tolerate therapeutic trials fo muscle relaxants, unless contraindicated, including but not limited to baclofen, cyclobenzaprine, orphenadrine, Skelaxin, and tizanidine.</p> <p>Migraine: Consideration for Botox approvals will only be made after failures of required trials of the following preferred medications: tricyclic or venlafaxine,beta blocker, valproic acid ,topiramate.</p> <p>Firdapse is recommended for the treatment of Lambert-Eaton myasthenic syndrome (LEMS) in adults.</p> <p>Ruzurgi is recommended for the treatment of Lambert-Eaton myasthenic syndrome (LEMS) in patients 6 years to less than 17 years of age.</p>
NEUROLOGICS- hATTR AGENTS				MC MC/DEL MC/DEL MC/DEL MC/DEL	AMVUTTRA ¹ ONPATTRO ¹ TEGSEDI ¹ VYNDAMAX ¹ VYNDAQEL ¹ WAINUA ¹	<p>1. PA required for appropriate diagnosis.</p> <p>Use PA Form# 20420</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.</p> <p>Tegsedi® should be non-preferred and approved for patients for whom other treatments, including Onpattro®, have been ineffective.</p> <p>Vyndamax will be considered for the treatment of the cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) in adults to reduce cardiovascular mortality and cardiovascular-related hospitalization</p>
NEUROLOGICS- SMA	MC MC MC		<p>GENE</p> <p>ZOLGENSMA¹</p> <p>NON-GENE</p> <p>EVRYSDI^{1,2} SPINRAZA¹</p>			<p>1. Clinical PA is required to establish diagnosis and medical necessity</p> <p>2. For patients 2 months of age and older.</p> <p>Use PA Form# 20420</p>	<p>Zolgensma: The patient is less than 2 years of age AND The diagnosis is spinal muscular atrophy (SMA) AND The patient has bi-allelic mutations of the SMN1 gene AND The patient does not have advanced SMA (e.g. complete paralysis of limbs or permanent ventilator dependence) AND Medication is prescribed per the dosing</p> <p>Spinraza: The diagnosis is spinal muscular atrophy (SMA) type 1, 2, or 3 (results of genetic testing must be submitted) AND The patient has at least 2 copies of the SMN2 gene AND The prescriber is a neurologist, pulmonologist, or other physician with expertise in treating SMA AND Baseline motor ability has been established using one of the following exams: Hammersmith Infant Neurological Exam (HINE) Hammersmith Functional Motor Scale Expanded (HFMSE) Upper Limb Module Test (non-ambulatory) Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND) AND Prior to starting therapy, and prior to each dose, the following laboratory tests will be conducted: Treating provider attests the member has a platelet count > 50,000/ml or greater Treating provider agrees to do platelet count and coagulation test before each dose Treating provider agrees to do a quantitative spot urine protein test before each dose Concomitant use of Spinraza and Zolgensma is investigational and will not be approved AND Use of Spinraza after gene replacement therapy, including Zolgensma is investigational and will not be approved Note: Initial approval will be granted for 4 loading doses (the first 3 loading doses should be administered at 14-day intervals; the 4th loading dose should be administered 30 days after the 3rd dose). Renewal may be granted for up to 12 months with a maximum of 3 doses approved per year (12mg (5ml) every 4 months). For therapy continuation, clinical documentation must be submitted documenting improvement or maintenance of motor ability OR slower progression of disease than would otherwise be expected.</p>
NEUROLOGICS- RETT SUNDROME				MC	DAYBUE ^{1,2}	<p>1. Clinical PA required for appropriate diagnosis</p> <p>2. For the treatment of patients 2 years of age and older.</p> <p>Use PA Form# 20420</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>

STEROIDS

GLUCOCORTICOIDS/ MINERALOCORTICOIDS	MC/DEL		BUDESONIDE EC 3mg DR CAPS	MC		ALKINDI SPRINKLE	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC		CELESTONE SUSP	MC		CORTEF 10 and 20 TABS		
	MC/DEL		CORTEF 5	MC/DEL		FLORINEF TABS		
	MC/DEL		CORTISONE ACETATE TABS	MC		HEMADY		
	MC/DEL		DELTASONE TABS	MC/DEL		MEDROL TABS		
	MC/DEL		DEPO-MEDROL SUSP	MC		MEDROL DOSEPAK TABS		
	MC/DEL		DEXAMETHASONE	MC		MILLIPRED		
	MC		DEXPAK	MC		ORTIKOS		
	MC/DEL		FLUDROCORTISONE ACETATE TABS	MC		ORAPRED SOLN		
	MC/DEL		HYDROCORTISONE	MC		PEDIAPRED LIQD		
	MC		KENALOG	MC		PREDNISONE INTENSOL CONC		
	MC/DEL		METHYLPREDNISOLONE TABS	MC		STERAPRED TABS		
	MC/DEL		PREDNISOLONE	MC		ZILRETTA		
	MC/DEL		PREDNISONE					
MC/DEL		SOLU-CORTEF SOLR						
MC/DEL		SOLU-MEDROL SOLR						

DDI: All preferred steroids will require clinical PA for patients over 60 that are currently on fluoroquinolone therapy.

HORMONE REPLACEMENT THERAPIES

ANDROGENS / ANABOLICS	MC/DEL		ANDRODERM PT24	MC		ANADROL-50	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Additionally, laboratory evidence of a testosterone deficiency must be supplied. One of each dosage form should be tried (tablet, injection, and topical)
	MC/DEL		ANDROGEL 1%	MC		ANDRO LA 200 OIL		
	MC/DEL		ANDROGEL PUMP 1.62%	MC/DEL		ANDROGEL PACKETS 1.62%		
	MC/DEL		DANAZOL CAPS	MC		ANDROID CAPS		
	MC/DEL		TESTOSTERONE CYP	MC		AXIRON		
				MC		DELATESTRYL OIL		
				MC/DEL		DEPO-TESTOSTERONE OIL		
				MC		FORTESTA		
				MC		HALOTESTIN TABS		
				MC/DEL		JATENZO		
				MC/DEL		METHITEST TAB		
				MC/DEL		METHYLTESTOSTERONE CAP		
				MC/DEL		OXANDROLONE		
				MC/DEL		STRIANT MUC ER		
			MC		TESTIM			
			MC/DEL		TESTOSTERONE GEL PACKETS			
			MC/DEL		TESTOSTERONE SOL			
			MC		TESTRED CAPS			
			MC		TLANDO			
			MC/DEL		VOGELXO			
			MC/DEL		XYOSTED			

Oxandrolone: Weight gain (adjunctive therapy): Adjunctive therapy to promote weight gain after weight loss following extensive surgery, chronic infections, or severe trauma, and in some patients who, without definite pathophysiologic reasons, fail to gain or to maintain normal weight. Other indications included in manufacturer labeling: Adjunctive therapy to offset protein catabolism with prolonged corticosteroid administration. Requirement for documentation of weight loss over two readings- Patient has involuntary weight loss of more than 10% of total body weight in less than four months) and, BMI < 18.5 (Normal BMI = 18.5 to 24.9)

ESTROGENS - PATCHES / TOPICAL	MC		EVAMIST	MC/DEL	5	ESTRADIOL PTWK	Use PA Form# 20420	Approved for failures on multiple oral estrogen agents after 90 day trials or if unable to swallow any oral medication.
	MC/DEL		MINIVELLE PATCH	MC/DEL	8	DIVIGEL ¹		
				MC/DEL	8	CLIMARA PTWK		
				MC/DEL	8	ELESTRIN ¹		
				MC/DEL	8	MENOSTAR PATCH		
				MC/DEL	8	VIVELLE-DOT PTTW		

ESTROGENS - TABS	MC/DEL		ESTRADIOL	MC/DEL		ENJUVIA	Use PA Form# 20420	Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		PREMARIN TABS	MC/DEL		ESTRADIOL-NORETHINDRONE		
				MC/DEL		ESTRACE TABS		
				MC		ESTRATAB TABS		
				MC/DEL		MENEST TABS		
				MC/DEL		NORETHINDRON-ETHINYL		
			MC		ORTHO-EST TABS			

	MC MC MC/DEL MC/DEL MC	TRI-LO-ESTARYLLA TAB TRI-ESTARYLLA TRI-SPRINTEC TAB TRI-LO-SPRINTEC TRINESSA						If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.
CONTRACEPTIVES - MULTI-PHASIC COMBINATIONS			MC		NATAZIA		Use PA Form# 20420 Use PA Form# 20420	
VASOMOTOR SYMPTOMS AGENTS								
VASOMOTOR SYMPTOMS AGENTS			MC/DEL		VEOZAH		Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid concomitant use of Veozah with drugs that are weak, moderate or strong CYP1A2 inhibitors. Veozah: Approval requires at least one preferred Hormone Replacement Therapy (HRT) and two preferred non-hormonal therapies (i.e., SSRIs, SNRIs, gabapentin, pregabalin, clonidine).
DIABETES SUPPLIES								
DIABETIC - SUPPLIES		CONTINUOUS GLUCOSE MONITORING ^{1,2} DIABETIC- LANCETS DIABETIC- LANCING DEVICES DIABETIC- LANCING DEVICES DIABETIC- PEN NEEDLES DIABETIC- SYRINGES DIABETIC- TEST STRIPS DIABETIC- METERS					Use PA Form#20420	1. Clinical PA is required to establish diagnosis and medical necessity. 2. Dosing limits apply. Please refer to Dose consolidation list. Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepdl.org Continuous Glucose Monitoring Criteria: Patient has a diagnosis of Diabetes Mellitus AND Practitioner feels patient has sufficient training to use CGM • 2 years of age or older for Dexcom G6 and Dexcom G7, ≥ 14 years for Medtronic Guardian, or ≥ 4 years for Freestyle Libre 2. • At least one of the following are documented: o Hypoglycemic unawareness o Treated with insulin (at least 1X day) o Has history of problematic hypoglycemia with documentation of at least one recurrent level 2 hypoglycemic events, or 1 level 3 hypoglycemic event • Approval of non-preferred products will be limited to cases where the CGM is directly integrated with the patient's insulin pump. The make and model of pump must be documented on the prior authorization.
DIABETES THERAPIES								
DIABETIC - INSULIN	MC/DEL MC MC MC MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	APIDRA HUMALOG KWIKPEN INJ 100/ML HUMALOG JUNIOR KWIKPEN 100/ML HUMALOG MIX 75/25 HUMALOG 50/50 VIAL HUMULIN INJ 70/30 KWIKPEN HUMULIN INJ 70/30 HUMULIN R INJ U-500 INSULIN ASPART PROT MIX 70-30 INSULIN ASPART INSULIN LISPRO LANTUS SOLN LEVEMIR NOVOLOG NOVOLOG MIX NOVOLOG MIX 70/30 FLEXPEN	MC/DEL MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC		ADMELOG AFREZZA ¹ BASAGLAR FIASP HUMALOG KWIKPEN U-200 HUMULIN INJ 50/50 HUMULIN N INJ U-100 HUMULIN R U-100 LYUMJEV NOVOLIN RELION		Use PA Form# 20420	1. Not to be as a monotherapy. Obtain lab values of pulmonary function and recent smoking history 2. For the treatment of patients ≥3 years of age Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - PENFILLS	MC MC	HUMALOG MIX KWIK 50/50 HUMALOG MIX INJ 75/25 KWP	MC MC/DEL		APIDRA OPTICLIK PEN NOVOLIN 70/30 PEN			Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	<p>MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL</p>	<p>HUMALOG KWIK INJ 100/ML HUMALOG KWIK INJ 200/ML HUMULIN R U-500 KWP INSULIN ASPART PROT MIX 70-30 PEN INSULIN ASPART PEN INSULIN LISPRO KWIKPEN U-100 LANTUS SOLOSTAR LEVEMIR FLEXTOUCH LEVEMIR FLEXPEN NOVOLOG MIX PENFILL NOVOLOG PENFILL SOLN NOVOLOG FLEXPEN NOVOLOG MIX 70/30 VIAL TOUJEO MAX SOLOSTAR TOUJEO SOLOSTAR</p>	<p>MC MC/DEL</p>	<p>REZVOGLAR KWIKPEN TRESIBA</p>	<p>another drug and the preferred drug(s) exists.</p>	
DIABETIC - DPP- 4 ENZYME INHIBITOR	<p>MC/DEL MC/DEL</p>	<p>JANUVIA^{1,2} TRADJENTA²</p>	<p>MC/DEL MC/DEL MC/DEL MC</p>	<p>NESINA ONGLYZA² QTERN ZITUVIO</p>	<p>1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile.</p> <p>2. Dosing limits apply. Please refer to Dose consolidation list.</p> <p>Use PA Form# 20420</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Onglyza 5mg will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A4 inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin).</p>
DIABETIC - DPP- 4 ENZYME INHIBITOR-COMBO	<p>MC/DEL MC/DEL MC/DEL</p>	<p>JANUMET^{1,2} JANUMET XR^{1,2} JENTADUETO¹</p>	<p>MC/DEL MC/DEL MC MC/DEL</p>	<p>JENTADUETO XR KAZANO KOMBIGLYZE XR OSENI</p>	<p>1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile.</p> <p>2. Dosing limits apply. Please refer to Dose consolidation list.</p> <p>Use PA Form# 20420</p>	
DIABETIC - LANCET-LANCET DEVICE					<p>Use PA Form# 20420</p>	Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepdl.org
DIABETIC - SYRINGES-NEEDLES					<p>Use PA Form# 20420</p>	Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepdl.org
DIABETIC - OTHER			<p>MC/DEL MC</p>	<p>CYCLOSET SYMLIN</p>	<p>Use PA Form #20420 for all others</p>	
SGLT 2 INHIBITORS	<p>MC/DEL MC/DEL MC/DEL</p>	<p>FARXIGA INVOKANA¹ JARDIANCE</p>	<p>MC/DEL</p>	<p>STEGLATRO</p>	<p>1.Dosing limits apply please refer to Dose Consolidation List</p> <p>Use PA Form# 20420</p>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

SGLT 2 INHIBITOR COMBINATIONS	MC/DEL MC/DEL MC/DEL		INVOKAMET SYNJARDY XIGDOU XR	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		GLYXAMBI INVOKAMET XR SEGLUROMET STEGLUJAN SYNJARDY XR TRIJARDY XR		Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Glyxambi /Xigduo XR- Verify prior trials and failures or intolerance of preferred treatments from other diabetic categories Synjardy® XR is not recommended for patients with type 1 DM or for the treatment of diabetic ketoacidosis. Use PA Form# 20420
DIABETIC MONITOR	MC MC MC MC		ONE TOUCH ULTRA 2 KIT ONE TOUCH ULTRA MINI KIT TRUE METRIX TRUETRACK	MC MC MC MC MC MC MC MC		ACCUCHECK ASCENSIA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE INSULINX FREESTYLE LITE SYSTEM KIT ONE TOUCH ULTRA SMART KIT PRECISION XTRA METER PRODIGY	Use PA Form# 20420	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters.
DIABETIC TEST STRIPS	MC MC MC		ONE TOUCH ULTRA ¹ TRUE METRIX TRUETRACK	MC MC MC MC MC MC MC MC MC		ACCUCHECK ASCENSIA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE FREESTYLE LITE FREESTYLE INSULINX ONE TOUCH DELICA PRECISION XTRA PRODIGY	1. Only 50 ct & 100 ct package size. Use PA Form# 20420	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters.
INCRETIN MIMETIC	MC MC MC/DEL		BYETTA TRULICITY VICTOZA	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL	5 5 8 8 8 8	OZEMPIC RYBELSUS ADLYXIN BYDUREON BCISE MOUNJARO SOLIQUA XULTOPHY		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Soliqua must try both insulin and a preferred incretin mimetic and have a medical necessity for use that is not based on convenience or simply due to the fact that one injection is needed instead of two. Use PA Form# 20420
DIABETIC - ORAL SULFONYLUREAS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		CHLORPROPAMIDE TABS GLIMEPIRIDE GLIPIZIDE TABS GLIPIZIDE ER TABS GLYBURIDE MICRONIZED TABS GLYBURIDE TABS ¹ TOLAZAMIDE TABS TOLBUTAMIDE TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AMARYL TABS DIABETA TABS GLUCOTROL TABS GLUCOTROL XL TBCR GLYNASE TABS MICRONASE TABS	Use PA Form# 20420 1. Pa required for members ≥65. Glyburide has a greater risk of severe prolonged hypoglycemia in older adults.	Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: All sulfonylureas (except glyburide) will now be non-preferred and require prior authorization if it is currently being used with either ranitidine or cimetidine. DDI: Glimepiride will now be non-preferred and require prior authorization if it is currently being used with either fluconazole (except 150mg strength) or fluvoxamine. Amaryl is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either fluconazole or fluvoxamine.
DIABETIC -ORAL BIGUANIDES	MC/DEL		METFORMIN HCL TABS	MC		GLUCOPHAGE TABS	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered

	MC/DEL		METFORMIN ER	MC MC MC/DEL	GLUCOPHAGE XR TB24 FORTAMET METFORMIN ER OSMOTIC		on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - THIAZOL / BIGUANIDE COMBO				MC/DEL MC/DEL MC MC	ACTOPLUS MET ¹ ACTOPLUS MET XR AVANDARYL ¹ AVANDAMET TABS ¹	Use PA Form# 20420 1. Requires use of Actos, Metformin, or other preferred anti-diabetics.	DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - / THIAZOL	MC/DEL		PIOGLITAZONE HCL ¹	MC/DEL MC	ACTOS TABS ³ AVANDIA TABS ²	1. Pioglitazone HCL is non-preferred as monotherapy. Pioglitazone HCL is preferred if therapeutic doses of metformin, sulfonylurea or insulin are seen in members drug profile for at least 60 days within the past 18 months. 2. Current users of Avandia who have tried Actos will be able to continue use of Avandia. 3. Dosing limits apply please refer to Dose Consolidation List Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - ALPHAGLUCOSIDASE	MC/DEL			MC	PRECOSE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - SULFONYLUREA / BIGUANIDE	MC/DEL		GLYBURIDE/METFORMIN	MC MC MC/DEL	GLUCOVANCE TABS ¹ METAGLIP TABS ¹ DUETACT ²	1. Use individual ingredients. 2. Use Actos with generic glimepiride. Use PA Form# 20420	Approved for patients failing to achieve good diabetic control with maximal doses of individual components.
DIABETIC - MEGLITINIDES	MC		NATEGLINIDE	MC/DEL MC/DEL	PRANDIN TABS STARLIX TABS	Use PA Form# 20420	Preferred drugs from other diabetic sub-categories must be tried and failed due to lack of inadequate diabetic control or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Prandin is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with both Sporanox and gemfibrozil, due to a significant drug-drug interaction.
GLUCOSE ELEVATING AGENTS							
GLUCOSE ELEVATING AGENTS	MC/DEL	1	GLUCAGEN INJ. HYPOKIT ¹	MC	GLUCAGON DIAGNOSTIC KIT	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	2	BAQSIMI ^{2,4}	MC MC/DEL MC	GLUCAGEN DIAGNOSTIC KIT GVOKE ³ ZEGALOGUE ⁵	1. Dosing limits apply, please see dose consolidation list. 2. For the treatment of patients ≥ 4 years of age. 3. For the treatment of patients ≥ 2 years of age. 4. Baqsimi will require a step through Glucagen. 5. For the treatment of patients ≥ 6 years of age.	

THYROID							
THYROID EYE DISEASE				MC	TEPEZZA	Use PA Form# 20420	
THYROID HORMONES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ARMOUR THYROID TABS CYTOMEL TABS ERMEZA ¹ LEVOTHROID TABS LEVOTHYROXINE SODIUM TABS LEVOXYL TABS UNITHROID TABS	MC MC/DEL MC MC/DEL	LEVOTHYROXINE SODIUM SOLR LIOETHYRONINE SYNTHROID TABS THYQUIDITY	Use PA Form# 20420 1.Clinical PA is required to confirm diagnosis of dysphagia.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTITHYROID THERAPIES	MC/DEL MC/DEL		METHIMAZOLE TABS PROPYLTHIOURACIL TABS	MC/DEL	TAPAZOLE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CUSHING DISEASE AGENTS							
CUSHING DISEASE AGENTS				MC MC	ISTURISA ¹ RECORLEV	1. For the treatment of adult patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative. Use PA Form #20420	Recorlev® is associated with dose-related QT interval prolongation. QT interval prolongation may lead to life-threatening ventricular dysrhythmias such as Torsades de pointes.
OSTEOPOROSIS / BONE AGENTS							
OSTEOPOROSIS	MC/DEL		ALENDRONATE	MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC MC MC MC/DEL	ACTONEL TABS ARELIA SOLR BINOSTO BONIVA INJECTION KIT BONIVA TABS ^{2,4} CALCITONIN NS DUAVEE DIDRONEL TABS EVISTA TABS ¹ EVENTY ² FORTEO FORTICAL FOSAMAX TABS AND PLUS D ³ PROLIA SOHONOS ⁶ STRENSIQ ⁵ TYMLOS XGEVA ZOMETA	Use PA Form# 20420 1. Approval only requires failure of Alendronate. 2. Quantity limits apply, please see dosage consolidation list. 3. Please use Alendronate and Vitamin D. 4. Please use other preferred agents. 5. Obtain baseline ophthalmology exams and renal ultrasounds and then periodically during treatment 6. Clinical PA for indication required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Binosto use preferred generic alendronate tablets Evenity® should be limited to 12 monthly doses Sohonos: For the reduction in volume of new heterotopic ossification in adults and pediatric patients aged 8 years and older for females and 10 years and older for males with fibrodysplasia ossificans progressiva (FOP).
FIBROBLAST GROWTH FACTOR 23 INHIBITORS	MC		CRYSVITA ¹			1.Preferred for patients <21 years for the treatment of X-linked hypophosphatemia. Use PA Form #20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CALCIMIMETIC AGENTS							
CALCIMIMETIC AGENTS				MC MC	PARSABIV SENSIPAR	Use PA Form# 30115	For Sensipar baseline PTH, Ca, and phosphorous levels are required and initial approvals will be limited to 3 months. Subsequent approvals will require additional levels being done to assess changes. Will not approve if baseline Ca is less than 8.4.

Parsabiv is for the treatment of secondary hyperparathyroidism (HPT) in adults with chronic kidney disease (CKD) on hemodialysis. Parsabiv® has not been studied in adults with parathyroid carcinoma, primary hyperparathyroidism, or with chronic kidney disease who are not on hemodialysis and is not recommended for use in these populations.

GROWTH HORMONE

GROWTH HORMONE	MC/DEL	GENOTROPIN ¹ NORDITROPIN SOLN ¹ NUTROPIN AQ ¹	MC	8	HUMATROPE SOLR	Use PA Form# 10710 1. Clinical PA is required to establish diagnosis and medical necessity.	See Growth Hormone PA form for criteria. Step-order will still apply unless clinical contraindication supplied. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		MC	8	INCRELEX		
			MC	8	NUTROPIN		
	MC/DEL		MC/DEL	8	NGENLA		
			MC	8	OMNITROPE		
			MC	8	SAIZEN SOLR		
			MC	8	SKYTROFA		
			MC/DEL	8	SOGROYA		
	MC/DEL	8	TEV-TROPIN				

ACHONDROPLASIA TREATMENT			MC		VOXZOGO ¹	1. Pediatric patients with achondroplasia who are 5 years of age and older with open epiphyses. Use PA Form# 20420	Voxzogo: To increase linear growth in pediatric patients with achondroplasia who are 5 years of age and older with open epiphyses. This indication is approved under accelerated approval based on an improvement in annualized growth velocity. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).
--------------------------	--	--	----	--	----------------------	---	---

SOMATOSTATIC AGENTS			MC/DEL	7	OCTREOTIDE INJ ¹	Use PA Form# 10710 1. Non-preferred products must be used in specified step order.	
			MC	8	BYNFEZIA ¹		
			MC	8	MYCAPSSA ¹		
			MC/DEL	8	SANDOSTATIN ¹		
			MC	8	SOMATULINE ¹		

GROWTH HORMONE ANTAGONISTS

GH ANTAGONISTS			MC		SOMAVERT	Use PA Form# 10710	Approved for acromegaly patients failing surgery/radiation/drug therapy including bromocriptine and sandostatin.
----------------	--	--	----	--	----------	------------------------------------	--

VASOPRESSIN RECEPTOR ANTAGONIST

VASOPRESSIN RECEPTOR ANTAGONIST			MC		JYNARQUE ¹	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis	Samsca Drug Warning- Avoid use in patients with underlying liver disease, including cirrhosis, because the ability to recover from liver injury may be impaired. Limit duration of therapy to 30 days to minimize the risk of liver injury. DDI: Jynarque- Concomitant use with strong CYP3A inhibitors is contraindicated. Avoid concomitant use of Jynarque® with OATP1B1/B3 and OAT3 substrates (e.g. statins, bosentan, glyburide, nateglinide, repaglinide, methotrexate, furosemide).
			MC/DEL		SAMSCA		

URINARY INCONTINENCE

VASOPRESSINS	MC/DEL	DESMOPRESSIN TABS DDAVP SOLN	MC/DEL	5	DDAVP TABS	1. Products must be used in specified step order. Nocturnal enuresis patients will be encouraged to periodically attempt stopping DDAVP. 2. Patients with a diagnosis of hemophilia or Von Willebrands disease will be exempt from prior authorization. Use PA Form# 20420	Approved for central diabetes insipidus and for nocturnal enuresis. For nocturnal enuresis- must be over 6 years old, must fail an adequate trial of alarm training (higher success rate, lower relapse rate) and must periodically attempt weaning (at 6 month intervals).
	MC/DEL		MC/DEL	6	DESMOPRESSIN SPRAY ¹		
			MC	8	DESMOPRESSIN ACETATE SOLN ¹		
			MC/DEL	8	NOCDURNA ¹		
			MC	8	NOCTIVA ¹		
			MC/DEL	8	STIMATE SOLN ^{1,2}		

ANTISPASMODICS	MC/DEL	DETROL TABS DETROL LA CAPS OXYBUTYNIN	MC/DEL	8	DARIFENACIN ER TAB	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		8	DITROPAN			
	MC/DEL		8	FLAVOXATE HCL TAB			
			8	TOLTERODINE			

ANTISPASMODICS - LONG ACTING	MC/DEL	GELNIQUE GEL PACKET	MC	8	DITROPAN XL TBCR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered
------------------------------	--------	---------------------	----	---	------------------	------------------------------------	--

	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	MYRBETRIQ OXYBUTYININ ER TABS OXYTROL SOLIFENACIN SUCCINATE TAB TOVIAZ TROSPIUM	MC/DEL MC MC/DEL MC MC MC	8 8 8 8 8 8	ENABLEX ^{1,2} GEMTESA ² TOLTERODINE TAB VESICARE ¹ VESICARE ³ LS	1. See Criteria Section. 2. Use a preferred long acting antispasmodic. 3. For the treatment of patients ≥ 2 years of age.	on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. 1. Vesicare 5mg and Enablex 7.5mg maximum doses if given with drugs known to be significant CYP3A4 inhibitors.(Ketoconazole, Sporanox, Erythromycin, Fluconazole, Nefazodone, Nelfinavir, and Ritonavir) DDI: Enablex 15mg and Vesicare 10mg will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: clarithromycin, erythromycin, Ketek, Crixivan, Norvir, ketoconazole, fluconazole (except 150mg strength), Sporanox, nefazodone, or diltiazem.
CHOLINERGIC	MC/DEL	BETHANECHOL	MC/DEL		URECHOLINE	Use PA Form# 20420	
HYPERAMMONIA TREATMENTS	MC	CARGLUMIC ACID TABS	MC		CARBAGLU TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
UREA CYCLE DISORDER	MC MC	BUPHENYL TABLET PHEBURANE GRANULES	MC MC MC MC/DEL MC/DEL		BUPHENYL POWDER RAVICTI LIQUID OLPRUVA SODIUM PHENYL BUTYRATE POWDER SODIUM PHENYL BUTYRATE TAB	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Olpruva: As adjunctive therapy to standard of care, which includes dietary management, for the chronic management of adult and pediatric patients weighing 20kg or greater and with a body surface area (BSA) of 1.2m2 or greater, with urea cycle disorders (UCDs) involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS).
METABOLIC MODIFIER							
HERED. TYROSINEMIA			MC		ORFADIN	Use PA Form# 20420	Approved for Type 1 hereditary tyrosinemia patients. Must include laboratory evidence of dx at first PA.
FABRY DISEASE AGENTS			MC MC MC/DEL		ELFABRIO ¹ FABRAZYME ² GALAFOLD ¹	Use PA Form# 20420	1.Clinical PA to verify appropriate diagnosis. 2.For the treatment of patients 2 years of age and older. Elfabrio and Galfold: For the treatment of adults with confirmed Fabry disease.
ANTIHYPERTENSIVES / CARDIAC							
CARDIAC GLYCOSIDES	MC/DEL MC/DEL MC/DEL	DIGITEK TABS DIGOXIN LANOXIN				Use PA Form# 20420	
CARDIAC MYOSIN INHIBITORS			MC		CAMZYOS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Camzyos: For the treatment of adults with symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) to improve functional capacity and symptoms. DDI: Concomitant use of Camzyos® with a moderate to strong CYP2C19 inhibitor or a strong CYP3A4 inhibitor is contraindicated.
CARDIAC - SINUS NODE INHIBITORS			MC		CORLANOR	Use PA Form#20420	In patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction ≤35%, who are in sinus rhythm with resting heart rate ≥70 beats per minute (bpm) and
CARDIAC - SOLUBLE GUANYLATE CYCLASE STIMULATORS			MC/DEL		VERQUVO		

						Use PA Form# 20420	
CARDIAC- SODIUM- GLUCOSE COTRANSPORTER 2 (SGLT2) INHIBITOR				MC		INPEFA ¹	1. To reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit in adults with: Heart failure or Type 2 diabetes mellitus, chronic kidney disease, and other cardiovascular risk factors. Other Preferred SGLT inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIANGINALS--Isosorbide Di-nitrate/ Mono-Nitrates	MC/DEL MC/DEL		ISOSORBIDE MONONITRATE TABS ISOSORBIDE MONONITRATE ER	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		DILATRATE SR CPCR ISORDIL TABS ISORDIL TITRADOSE TABS ISOSORBIDE DINITRATE SUBL ISOSORBIDE DINITRATE TABS ISOSORBIDE DINITRATE CR TBCR ISOSORBIDE DINITRATE ER TBCR ISOSORBIDE DINITRATE TD TBCR IMDUR TB24 ISMO TABS MONOKET TABS	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NITRO - OINTMENT/CAP/CR	MC/DEL MC/DEL MC MC		NITROBID OINT NITROGLYCERIN CPCR NITROL OINT NITRO-TIME CPCR				Use PA Form# 20420
NITRO - PATCHES	MC/DEL MC/DEL	1 1	NITROGLYCERIN PT24 ¹ NITRO-DUR PT 24 0.8MG ¹	MC MC/DEL		NITRODISC PT24 NITRO-DUR PT24	1. At least 2 step 1's and step 3 of the preferred products must be used in specified order or PA will be required. Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NITRO - SUBLINGUAL/ SPRAY	MC/DEL		NITROSTAT SUBL	MC/DEL MC MC		NITROQUICK SUBL NITROLINGUAL SOLN NITROLINGUAL TABS	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS - NON SELECTIVE	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		CARVEDILOL LEVATOL TABS NADOLOL TABS PINDOLOL TABS PROPRANOLOL HCL SOLN ¹ PROPRANOLOL HCL TABS ¹ PROPRANOLOL HCL 60MG TABS PROPRANOLOL LA CAPS RANOLAZINE ER TABS SOTALOL AF SOTALOL HCL TABS TIMOLOL MALEATE TABS	MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC		ASPRUZYO BETAPACE TABS BETAPACE AF TABS COREG CR ³ COREG TABS CORGARD TABS INDERAL TABS HEMANGEOL SOL INDERAL XL CAP INDERAL LA CPCR INNOPRAN XL RANEXA	1. Recommend using BID since its effects do not last 24 hours. 2. Please use other strengths in combination to obtain this dose. 3. Dosing limits still apply. Please see dose consolidation list Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Concomitant use of Ranolazine products with strong CYP3A inhibitors, including ketoconazole, itraconazole, clarithromycin, nefazodone, nelfinavir, ritonavir, indinavir, and saquinavir, is contraindicated.
BETA BLOCKERS - CARDIO SELECTIVE	MC/DEL MC/DEL MC/DEL		ACEBUTOLOL HCL CAPS ATENOLOL TABS ¹ BETAXOLOL HCL TABS	MC MC/DEL MC		KERLONE TABS LOPRESSOR TABS SECTRAL CAPS	1. Recommend using Atenolol (and metoprolol) BID since its effects do not last 24 hours Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		BISOPROLOL FUMARATE TABS BYSTOLIC METOPROLOL TARTRATE TABS ¹ METOPROLOL ER NEBIVOLOL HCL TAB	MC/DEL MC/DEL MC/DEL		TENORMIN TABS TOPROL XL TB24 ZEBETA TABS		Use PA Form# 20420		
BETA BLOCKERS - ALPHA / BETA	MC/DEL		LABETALOL HCL TABS	MC		TRANDATE TABS		Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
BETA BLOCKERS & DURECTIC COMBOS	MC/DEL		METOPROLOL-HYDROCHLOROTHIAZIDE TAB	MC/DEL		DUTOPROL		Use PA Form# 20420		
CALCIUM CHANNEL BLOCKERS-- Amlodipines, Bepridil, Diltiazems, Felodipines, Isradipines, Nifedipines, Nisoldipine, and Verapamils	MC/DEL		AMLODIPINE ¹	MC/DEL MC MC/DEL		KATERZIA NORLIQVA NORVASC TABS ¹		Use PA Form# 20420	1. Dosing limits apply, please see dose consolidation list.	
	MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		DILTIA XT CP24 DILTIAZEM HCL ER CP24 DILTIAZEM HCL XR CP24 DILTIAZEM CD 300MG CP24 DILTIAZEM CD 360MG CP24 CARTIA XT CP24 ¹ DILTIAZEM CD CP24 ¹ DILTIAZEM HCL ER CP24 ¹ DILTIAZEM XR CP24 ¹ TIAZAC CP24 ¹	MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL	5 6 8 8 8 8 8 8 8	DILACOR XR CP24 ¹ TAZTIA ¹ CARDIZEM TABS ¹ CARDIZEM CD CP24 ¹ CARDIZEM LA TB24 ¹ CARDIZEM SR CP12 ¹ DILTIAZEM HCL TABS ¹ DILTIAZEM HCL ER CP12 ¹ DILTIAZEM HCL ER CP12 ¹		Use PA Form# 20420	1. Products must be used in specified order or PA will be required. Just write "Diltiazem 24-hour" and the pharmacy will use a preferred long acting diltiazem that does not require PA.	
				MC/DEL MC/DEL		PLENDIL TB24 FELODIPINE		Use PA Form# 20420	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
				MC MC		DYNACIRC CAPS DYNACIRC CR TBCR ¹		Use PA Form# 20420	1. Established users will be grandfathered	
				MC MC		CARDENE SR CPR NICARDIPINE HCL CAPS		Use PA Form# 20420	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		AFEDITAB CR NIFEDIAC CC NIFEDICAL XL TBCR NIFEDIPINE TBCR NIFEDIPINE ER TBCR	MC/DEL MC/DEL MC/DEL MC/DEL		ADALAT CC TBCR ¹ NIFEDIPINE CAPS PROCARDIA CAPS PROCARDIA XL TBCR		Use PA Form# 20420	1. Established users of Adalat CC are grandfathered.	
				MC MC		SULAR TB24 SULAR CR ¹		Use PA Form# 20420	1. Established users of 10MG and 20MG strengths are grandfathered.	
	MC/DEL MC/DEL MC/DEL	1 1 1	VERAPAMIL HCL CR TBCR VERAPAMIL HCL ER TBCR VERAPAMIL HCL SR TBCR	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		CALAN TABS CALAN SR TBCR COVERA-HS TBCR ISOPTIN-SR VERAPAMIL HCL ER CP24 VERAPAMIL HCL SR CP24 VERAPAMIL HCL TABS VERELAN CP24 VERELAN PM CP24		Use PA Form# 20420	Products must be used in specified order or PA will be required. Just write "Verapamil 24-hour" and the pharmacy will use a preferred long acting generic that does not require PA.	
	ANTIARRHYTHMICS	MC/DEL MC/DEL MC/DEL MC/DEL		AMIODARONE HCL DISOPYRAMIDE FLECAINIDE MEXILETINE HCL	MC/DEL MC/DEL MC/DEL MC/DEL		CORDARONE DISOPYRAMIDE MULTAQ NORPACE		1. Prescription must be written by Cardiologist.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC/DEL MC/DEL MC MC/DEL MC/DEL		PROCAINAMIDE PROPAFENONE QUINAGLUTE QUINIDINE GLUCONATE QUINIDINE SULFATE	MC/DEL MC MC/DEL MC/DEL MC MC/DEL		PACERONE QUINIDEX TAMBOCOR TIKOSYN ¹ RYTHMOL SR RYTHMOL	Use PA Form# 20420	DDI: Amiodarone will now be non-preferred and require prior authorization if it is currently being used in combination with either Lovastatin (doses greater than 40mg/day) or Lipitor (doses greater than 20mg/day) or Levofloxacin or Gemifloxacin, or Moxifloxacin, or Ofloxacin. DDI: Multaq will be preferred unless the following medications are seen in the member's drug profile within the last 35 days for brand name medications or 90 days for generic medications: Erythromycin, Amiodarone and other antiarrhythmics, TCA's, Phenothiazine, Ketoconazole, Itraconazole, Voriconazole, Cyclosporine, Telithromycin, Clarithromycin, Nefazodone, Ritonavir.
ACE INHIBITORS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		BENAZEPRIL HCL CAPTOPRIL TABS ENALAPRIL MALEATE TABS FOSINOPRIL SODIUM LISINOPRIL TABS RAMIPRIL QUINAPRIL HCL	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL	5 5 8 8 8 8 8 8 8 8 8 8 8 8	MAVIK TABS ACCUPRIL TABS ACEON TABS ¹ ALTACE CAPS ¹ EPANED LOTENSIN TABS ¹ MOEXIPRIL HCL ¹ MONOPRIL HCT TABS ¹ PRINIVIL TABS ¹ QBRELIS UNIVASC ¹ VASOTEC TABS ¹ ZESTRIL TABS ¹	1. Non-preferred products must be used in specified order. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Non-preferred products are subject to step-order requirements unless clinical circumstances warrant exception.
ANGIOTENSIN RECEPTOR BLOCKER	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		AMLODIPINE-OLMESARTAN TAB ³ IRBESARTAN ¹ LOSARTAN ¹ MICARDIS TABS ³ OLMESARTAN ¹ TELMISARTAN ¹	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	8 8 8 8 8 8	ATACAND TABS AVAPRO BENICAR TABS COZAAR DIOVAN EDARBI TEVETEN TABS	Use PA Form# 20420 1. Dosing limits apply, please see dose consolidation list. 2. Use preferred active ingredients which are available without PA. 3. Preferred without a PA only if patient on a diabetic therapy or prior ACE therapy.	Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy
DIRECT RENIN INHIBITOR				MC/DEL MC/DEL MC/DEL		AMTURNIDE TEKTURNA ¹ TEKAMLO	1. Must show failure of single and combination therapy from all preferred antihypertensive categories. Use PA Form# 20420	
ANTIHYPERTENSIVES - CENTRAL	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		CLONIDINE HCL TABS GUANFACINE HCL TABS HYDRALAZINE HCL TABS HYLOREL TABS METHYLDOPA TABS MINOXIDIL TABS PRAZOSIN HCL CAPS RESERPINE TABS	MC/DEL MC/DEL MC MC MC/DEL MC MC/DEL		CLONIDINE PATCH CLONIDINE TTS GUANABENZ ACETATE TABS ISMELIN TABS MINIPRESS CAPS NEXICLON TENEX TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ACE INHIBITORS AND CA CHANNEL BLOCKERS				MC/DEL MC MC MC/DEL	8 8 8 9	AMLODIPINE/BENAZEPRIL PRESTALIA ¹ TARKA TBCR LOTREL CAPS	1. Prestalia will only be approved for patients ≥ 18 years of age. Use individual preferred generic medications. Use PA Form# 20420	
ACE AND THIAZIDE COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		BENAZEPRIL HCL/HYDROCHLOR CAPTOPRIL/HYDROCHLOROTHIA ENALAPRIL MALEATE/HCTZ TABS LISINOPRIL-HCTZ TABS LOTENSIN HCT TABS	MC/DEL MC MC/DEL MC/DEL MC MC/DEL		ACCURETIC TABS MONOPRIL HCT TABS PRINZIDE TABS UNIRETIC TABS VASERETIC TABS ZESTORETIC TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

BETA BLOCKERS AND DIURETIC COMBO'S	MC/DEL MC/DEL MC/DEL		ATENOLOL/CHLORTHALIDONE BISOPROLOL FUMARATE/HCTZ PROPRANOLOL/HCTZ	MC/DEL MC/DEL MC MC MC/DEL		CORZIDE TABS LOPRESSOR HCT TABS TENORETIC TIMOLIDE 10/25 TABS ZIAC TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ARB'S AND CA CHANNEL BLOCKERS	MC/DEL MC/DEL MC/DEL		AMLODIPINE/VALSARTAN AMLODIPINE/VALSARTAN HCT TRIBENZOR	MC/DEL MC MC/DEL MC/DEL		AZOR BYVALSON EXFORGE EXFORGE HCT	Use PA Form# 20420	DDI: Byvalson will be non-preferred and require a prior authorization if it is currently being used in combination with drugs known to be significant CYP2D6 inhibitors (e.g. quinidine, propafenone, fluoxetine, paroxetine). Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy
ARB'S AND DIURETICS	MC/DEL MC/DEL MC/DEL MC/DEL		BENICAR HCT ¹ LOSARTAN HCT ¹ MICARDIS HCT TABS ¹ VALSARTAN-HCT ¹	MC/DEL MC/DEL MC MC/DEL MC/DEL MC	7 8 8 8 8 8	IRBESARTAN HYDROCHLOROTHIAZIDE ATACAND HCT TABS AVALIDE TABS ¹ DIOVAN HCT TABS ¹ HYZAAR TABS TEVETEN HCT TABS	Use PA Form# 20420	1. Dosing limits apply, please see dose consolidation list. Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy
ANGIOTENSIN MODULATORS-ARB COMBINATION	MC		ENTRESTO	MC/DEL MC		EDARBYCLOR ENTRESTO SPRINKLES	Use PA Form# 20420	
ARB'S AND DIRECT RENIN INHIBITOR COMBINATION				MC/DEL		VALTURNA	Use PA Form# 20420	
DIURETICS	MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ACETAZOLAMIDE TABS BUMETANIDE CHLOROTHIAZIDE TABS CHLORTHALIDONE TABS EDECIN TABS EDECIN TABS HYDROCHLOROTHIAZIDE INDAPAMIDE TABS METHAZOLAMIDE TABS METHYCLOTHIAZIDE TABS SPIRONOLACTONE SPIRONOLACTONE/HYDRO TORSEMIDE TABS TRIAMTERENE/HCTZ ZAROXOLYN TABS	MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ALDACTAZIDE TABS ALDACTONE TABS AMILORIDE HCL BUMEX TABS DEMADEX TABS DIAMOX DIURIL DYAZIDE CAPS CAROSPIR ENDURON TABS FUROSCIX INSPIRA KERENDIA KEVEYIS LASIX TABS MAXZIDE MICROZIDE CAPS MIDAMOR TABS NAQUA TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Furoscix: The indication for use is the treatment of congestion due to fluid overload in adults with NYHA Class II or Class III chronic heart failure AND the medication is being prescribed by or in consultation with a cardiologist AND the patient is experiencing symptoms despite compliance with oral loop diuretic therapy AND oral loop diuretic therapy will be resumed as soon as practical AND medical reasoning beyond convenience is provided for not pursuing therapy in an outpatient infusion setting. PA approval will be authorized for 1 month. DDI: The concomitant use of Keveyis® with high dose aspirin is contraindicated.
CCB / LIPID				MC/DEL		CADUET	Use PA Form# 20420	
NEUROGENIC ORTHOSTATIC HYPOTENSION								
NEUROGENIC ORTHOSTATIC HYPOTENSION				MC		NORTHERA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
LIPID DRUGS								
CHOLESTEROL - BILE SEQUESTRANTS	MC/DEL MC/DEL		CHOLESTYRAMINE COLESTIPOL HCl	MC/DEL MC/DEL MC MC/DEL		COLESTID PREVALITE QUESTRAN WELCHOL TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CHOLESTEROL - FIBRIC ACID DERIVATIVES	MC/DEL MC/DEL MC/DEL		FENOFIBRATE TAB GEMFIBROZIL TABS NIACIN ER	MC MC/DEL MC/DEL MC/DEL		ANTARA LOPID FENOFIBRATE 120mg TAB FENOFIBRATE CAP FIBRICOR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Fenofibrate is preferred but will require a prior authorization requests if used concurrent with Warfarin.

Additional criteria for the diagnosis of homozygous familial hypercholesterolemia (Repatha only): Total cholesterol levels > 290mg/dL or LDL-C > 190mg/dL (adults) OR Total cholesterol levels > 260mg/dL or LDL-C > 155mg/dL (children < 16 years) and TG within reference range OR Confirmation of diagnosis by gene testing.

[Use PA Form# 20420](#)

PULMONARY ANTI-HYPERTENSIVES

PULMONARY ANTI-HYPERTENSIVES	<p>MC MC/DEL MC/DEL MC</p>		<p>EPOPROSTENOL INJ^{3,6} SILDENAFIL TADALAFIL VENTAVIS³</p>	<p>MC/DEL MC MC/DEL MC MC MC MC MC MC MC/DEL MC MC MC MC MC/DEL</p>		<p>ADEMPAS^{1,3} ADCIRCA⁴ ALYQ TAB FLOLAN³ LIQREV OPSUMIT^{1,2} OPSYNVI⁴ ORENITRAM REMODULIN³ REVATIO⁴ TADLIQ⁴ TYVASO UPTRAVI VELVETRI³ WINREVAIR⁴</p>	<p>1. Requires previous trials/failure of multiple preferred medications. 2. Dosing limits apply, please see the dose consolidation list. 3. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 3 or 4. 4. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA (WHO) functional class 2 or 3.</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Sildenafil will be preferred with clinical PA for treatment of pulmonary arterial hypotension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of Sildenafil with moderate or strong Cyp3A inhibitors DDI: Upravi will require a prior authorization if it is currently being used in combination with strong inhibitors of CYP2C8 (gemfibrozil) DDI: Opsumit will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin). DDI: Adempas will require a prior authorization if it is currently being used in combination with drugs known to be PDE inhibitors should be avoided (including dipyridamole, addira and tadalafil) with adempas Liqrev: treatment of pulmonary arterial hypertension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of Liqrev with moderate or strong CYP3A inhibitors.</p>
------------------------------	--	--	---	--	--	--	---	---

[Use PA Form# 20420](#)

ERA / ENDOTHELIN RECEPTOR ANTAGONIST	<p>MC MC</p>		<p>LETAIRIS^{1,2} TRACLEER</p>				<p>1. Providers must be registered with LEAP Prescribing program, a restricted distribution program. 2. Clinical PA is required to establish diagnosis and medical necessity.</p>	<p>Tracleer approvals will require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 2 thru 4. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer. Letairis approvals will require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and functional class 2 or 3 symptoms.</p>
--------------------------------------	--------------------------------	--	--	--	--	--	--	--

[Use PA Form# 20420](#)

IMPOTENCE AGENTS

IMPOTENCE AGENTS							<p>As of January 1, 2006, per CMS (federal govt.), impotence agents are no longer covered.</p>	<p>As of January 1, 2006, per CMS (federal govt.), impotence agents are no longer covered.</p>
------------------	--	--	--	--	--	--	--	--

ANTI-EMETOGENICS

ANTIEMETIC - ANTICHOLINERGIC / DOPAMINERGIC	<p>MC MC/DEL MC MC/DEL MC</p>		<p>BONJESTA MECLIZINE HCL TABS PROMETHAZINE SUPP PROMETHAZINE TRANSDERM-SCOP PT72</p>	<p>MC MC MC MC MC</p>		<p>ANTIVERT TABS BARHEMSYS PHENERGAN SOLN PROMETHAZINE 50MG SUPP PROMETHEGAN SUPP TORECAN TABS</p>	<p>Use PA Form# 20420</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Concomitant use of MAOIs and Bonjesta® is contraindicated.</p>
ANTIEMETIC - 5-HT3 RECEPTOR ANTAGONISTS/ SUBSTANCE P NEUROKININ	<p>MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL</p>		<p>DICLEGIS DRONABINOL CAPS GRANISETRON TAB ONDANSETRON TAB ONDANSETRON ODT TBDP ONDANSETRON SOL</p>	<p>MC MC MC MC MC MC</p>	<p>8 8 8 8 8 8</p>	<p>AKYNZEO¹ APREPITANT ALOXI ANZEMET TABS APONVIE⁴ CESAMET¹ CINVANTI⁴</p>	<p>1. Approvals will require diagnosis of chemo-induced nausea/vomiting and failed trials of all preferred anti-emetics, including 5-HT3 class (Ondansetron) and Marinol.</p>	<p>Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. * Ondansetron limits still apply as listed on the Ondansetron PA form for covered indications including chemotherapy, radiotherapy, post operative nausea & vomiting and hyperemesis gravidarum. Other medical indications will be approved or denied on a case by case basis. Hyperemesis and other medical indications approved are still subject to failure of multiple preferred antiemesis drugs.</p>

				MC	8	EMEND ²		Akynzeo- Concomitant use should be avoided in patients who are chronically using a strong CYP3A inducer such as rifampin.
				MC	8	FOCINVEZ ^{1,2}		
				MC/DEL	8	KYTRIL	2. Clinical PA is required for members on highly emetic anti-neoplastic agents.	Varubi – Available to the few who are unable to tolerate or who have failed on preferred medications
				MC/DEL	8	MARINOL CAPS		
				MC	8	SANCUSO		Aponvie is for the prevention of postoperative nausea and vomiting (PONV) in adults.
				MC	8	SUSTOL		
				MC	8	SYNDROS	3. Dosing limits apply, please see Dosage Consolidation List	
				MC	8	TRIMETHOBENZAMIDE CAP		
				MC	8	VARUBI		
				MC/DEL	8	ZOFRAN ODT TBDP ³	4. Clinical PA required for appropriate diagnosis	
				MC/DEL	8	ZOFRAN TABS ³		
				MC/DEL	8	ZOFRAN INJ ³		
				MC	8	ZUPLENZ		
							Use PA Form# 20420	
NON-SEDATING ANTIHISTAMINES / DECONGESTANTS								
ANTIHISTIMINES - NON-SEDATING	MC MC/DEL MC/DEL MC		ALAVERT TABS CETIRIZINE TABS LORATADINE TAVIST ND (OTC)	MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL	5 5 5 5 8 8 8 8 8 9	CLARINEX TABS ^{1,5} CLARINEX SYR ^{1,2} FEXOFENADINE ¹ ZYRTEC ¹ ZYRTEC SYR ^{1,2} ALLEGRA ³ CLARITIN ³ DESLORATADIN LORATADINE ODT ⁴ LEVOCETIRIZINE ⁴ XYZAL ³	1. Must fail preferred drugs, OTC loratidine and cetirizine before moving to non-preferred step order drugs. 2. Clarinex and Zyrtec syrup <6 yr w/o PA. 3. Must fail all step 5 drugs (Clarinex, Fexofenadine and Zyrtec) before moving to next step product. 4. All OTC versions of loratidine ODT are now non-preferred. 5. Pa's for Clarinex RediTabs will only be approved if between the ages of 6-11 years old.	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. No combination product with decongestant will be approved since pseudoephedrine available without PA. Pseudoephedrine is available with prescription.
							Use PA Form# 20530	
ANTIHISTIMINES - OTHER	MC/DEL MC/DEL MC/DEL		CLEMASTINE CHLORPHENIRAMINE DIPHENHYDRAMINE				Use PA Form# 20530	
ALLERGY / ASTHMA THERAPIES								
ANAPHYLACTIC DEVICES	MC/DEL MC/DEL MC/DEL		EPINEPHRINE EPIPEN EPIPEN JR	MC MC/DEL		TWINJECT SYMJEPI		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
							Use PA Form# 20420	
ALLERGEN IMMUNOTHERAPY				MC MC MC MC MC		ODACTRA ORALAIR ¹ PALFORZIA RAGWITEK GRASTEK	1. See criteria section	Prescriber must provide the testing to show that the patient is allergic to the components in the prescribed therapy and must provide a clinically valid rationale why single agent sublingual therapy is being chosen over subcutaneous therapy Palforzia® is approved for use in patients with a confirmed diagnosis of peanut allergy. Initial dose escalation may be administered to patients aged 4 through 17 years. Up-dosing and maintenance may be continued in patients 4 years of age and older. Odactra® is approved for use in persons 12 through 65 years of age. Note that Odactra® is not indicated for the immediate relief of allergic symptoms.
							Use PA Form# 20420	Treatment must start 12 weeks before expected onset of pollen season and only after confirmed by positive skin test or in vitro testing for pollen-specific IgE antibodies for any of the 5

							grass species contained in Oralair Oralair: Patient age ≥10 years and ≤65 years Have an auto-injectable epinephrine on-hand	
ANTIASTHMATIC - ANTICHOLINERGICS - INHALER	MC MC/DEL MC/DEL		INCRUSE ELLIPTA ³ SPIRIVA HANDIHALER ^{1,2} SPIRIVA RESPIMAT	MC/DEL MC MC/DEL		FLUTICASON-SALMETEROL LONHALA MAGNAIR TUDORZA	Use PA Form# 20420 1. Quantity limit of 1 inhalation daily (1 capsule) 2. We ask physicians to write "asthma" on the prescription whenever Spiriva is primarily being used for that condition. 3. Quantity limit of 1 inhalation daily	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - PHOSPHODIESTERASE 4 INHIBITORS	MC/DEL		ROFLUMILAST	MC/DEL MC		DALIRESP OHTUVAYRE ¹	Use PA Form# 20420 1. For the maintenance treatment of chronic obstructive pulmonary disease (COPD) in adult patients	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTICHOLINERGICS - NEBULIZER	MC/DEL		IPRATROPIUM BROMIDE SOLN	MC MC/DEL		ATROVENT SOLN YUPELRI	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTIINFLAMMATORY AGENTS	MC/DEL MC/DEL MC/DEL MC MC/DEL		CROMOLYN SODIUM NEBU DUPIXENT ^{2,4} FASENRA ² FASENRA ² AUTO INJCT NUCALA ² SYRINGE 40MG XOLAIR ¹	MC MC		CINQAIR ³ TEZSPIRE ⁵	1. Need max inhaled steroids and written by pulmonary or allergy specialist. Must have elevated IgE and ≥ to age 6. 2. For patients with severe asthma aged 12 years or older and eosinophilia. 3. For patients ≥ 18 years of age with eosinophilia. 4. Clinical PA required. 5. For adult and pediatric patients aged 12 years and older with severe asthma. Use PA Form# 20420	All will require suboptimal response to maximal doses of inhaled steroid as evidenced by asthmatic ER/Hospital admissions and Allergy/Pulmonary specialist management. Dupixent limited to patient with asthma not controlled on high dose ICS-LABA who have eosinophil greater than or equal to 150 cells or the patient is depend on an oral corticosteroid Fasenra, Nucala and Cinqair are not indicated for treatment of other eosinophilic conditions and are not indicated for the relief of acute bronchospasm or status asthmaticus.
ANTIASTHMATIC - NASAL STEROIDS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC		BUDESONIDE SPRAY FLUTICASON SPR ³ OLOPATADINE SPRAY OMNARIS SPR ³ TRIAMCINOLONE NS QNASL	MC MC/DEL MC/DEL MC/DEL MC/DEL MC	5 8 8 8 8 8	BECONASE AQ INHA ^{1,3} DYMISTA FLONASE SUSP ^{2,3} FLUNISOLIDE SOLN ^{1,3} NASONEX SUSP RHINOCORT AERO ^{2,3}	Use PA Form# 20420 1. All preferred drugs must be tried before moving to non preferred steps. 2. All step 5 medications	Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

				MC/DEL MC MC MC MC/DEL MC MC/DEL	8 8 8 8 8 8 8	RHINOCORT AQUA SUSP ^{2,3} RYALTRIS ⁴ TRI-NASAL SOLN ^{2,3} VANCENASE POCKETHALER AERS ^{2,3} VERAMYST ^{2,3} XHANCE ² ZETONNA ³	need to be tried before moving to step 8's. 3. Dosing limits apply to whole category, please see dosage consolidation list. 4. Use of individual ingredients or other preferred agents.	Xhance will be considered for the treatment of nasal polyps in patients 18 years of age or older. The patient has had a documented side effect, allergy, or treatment failure of two preferred nasal glucocorticoids, one of which must be fluticasone.
ANTIASTHMATIC - NASAL MISC.	MC/DEL MC/DEL MC		AZELASTINE CROMOLYN NASAL 4% IPRATROPIUM NASAL SOL ¹	MC/DEL MC/DEL	8 8	ASTEPRO ² PATANASE	Use PA Form# 20420 1. Ipratropium will be approved if submitted with documentation supporting use of CPAP machine. 2. Utilize Multiple preferred, as well as step therapy Azelastine.	Approved if patient fails on non-sedating antihistamines and steroid nasal sprays. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - BETA - ADRENERGICS	MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC		ALBUTEROL NEB METAPROTERENOL PROAIR RESPICLICK PROVENTIL HFA SEREVENT TERBUTALINE SULFATE TABS ALBUTEROL 0.63mg/3ml VENTOLIN HFA AERS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC MC		ACCUNEB NEBU ALBUTEROL HFA BRETHINE LEVALBUTEROL TARTRATE PROAIR DIGIHALER ⁴ STRIVERDI VOLMAX TBCR VOSPIRE ER TB12 XOPENEX HFA ³ XOPENEX NEBU ^{1,2}	1. Xopenex users w/ prior asthma hospitalization due to albuterol nebulizer failure will be grandfathered. 2. Quantity Limit: 12 cc/day. 3. Dosing limits apply, please see dosage consolidation list. 4. For the treatment of patients ≥ 4 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ADRENERGIC COMBINATIONS	MC MC MC MC MC/DEL MC/DEL		ADVAIR DISKUS ¹ ADVAIR HFA ¹ AIRDUO RESPICLICK ² BREQ ELLIPTA ¹ DULERA SYMBICORT	MC MC/DEL MC/DEL MC		AIRDUO DIGIHALER ² AIRSUPRA BREZTRI AEROSPHERE TRELEGY ELLIPTA ¹	1. Dosing limits apply, please see dosage consolidation list. 2. For patients ≥ 12 years and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. AirDuo® Respiclick be non-preferred and require prior authorization and be available to those who are unable to tolerate or who have failed on preferred medications DDI: Avoid concomitant use of strong CYP3A4 inhibitors (e.g. ritonavir, atazanavir, clarithromycin, indinavir, itraconazole, nefazodone, nelfinavir, saquinavir, ketoconazole, telithromycin) with AirDuo® Respiclick is not recommended due to increased systemic corticosteroid and increased cardiovascular adverse effects
ANTIASTHMATIC - ADRENERGIC ANTICHOLINERGIC	MC/DEL MC MC/DEL MC/DEL		ALBUTEROL/IPRATROPIUM NEB. SOLN ANORO ELLIPTA COMBIVENT RESPIMAT STIOLTO	MC/DEL MC/DEL MC/DEL		BEVESPI AEROSPHERE ^{2,3} DUAKLIR PRESSAIR DUONEB SOLN ¹	1. Please use preferred individual ingredients Albuterol and Ipratropium. 2. Dosing limits apply, please see dosing consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Duoneb components are available separately without PA. DDI: Avoid concomitant use of Bevespi with other anticholinergic-containing drugs, due to an increased risk of anticholinergic adverse events. Bevespi® should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval.

						3. The safety and efficacy of use in children under the age of 18 years have not been established. Use PA Form# 20420	Bevespi should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval.
ANTIASTHMATIC - XANTHINES	MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AMINOPHYLLINE TABS THEOCHRON TB12 THEOLAIR-SR TB12 THEOPHYLLINE CR TB12 THEOPHYLLINE ELIX THEOPHYLLINE SOLN THEOPHYLLINE ER CP12 THEOPHYLLINE ER TB12	MC/DEL MC MC/DEL		THEO-24 CP24 THEOLAIR TABS UNIPHYL TBCR	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - STEROID INHALANTS	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC		ARNUITY ELLIPTA ASMANEX TWISTHALER ^{3,4} ASMANEX HFA ⁵ BUDESONIDE NEB 0.25MG & 0.5MG ¹ FLOVENT DISKUS ³ PULMICORT FLEXHALER ³ QVAR AERS ³	MC MC/DEL MC MC/DEL MC/DEL MC	8 8 8 8 8 8	AEROSPAN ALVESCO ³ ARMONAIR DIGIHALER BUDESONIDE NEB 1MG PULMICORT SUSP FLOVENT HFA ³	1. Budesonide Neb 0.25mg & 0.5mg will be preferred for members under the age of 8 years old. PA will be required for members 8 years of age and older, please consider other preferred options. 2. All preferreds must be Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

							<p>tried before moving to non preferred steps.</p> <p>3. Dosing limits apply, please see dosage consolidation list.</p> <p>4. Asmanex 110mcg will be limited to member between the ages of 4-11years old.</p> <p>5. Asmanex HFA will be preferred for members under the age of 6 years old. PA will be required for members 6 years of age and older, please consider other preferred options.</p> <p>Use PA Form# 20420</p>	
ANTIASTHMATIC - 5-Lipoxygenase Inhibitors				MC		ZYFLO CR TABS		Other Preferred asthma controller drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - LEUKOTRIENE RECEPTOR ANTAGONISTS	MC/DEL MC/DEL MC/DEL		MONTELUKAST GRANULE ¹ MONTELUKAST SODIUM TAB MONTELUKAST SODIUM CHEW TAB	MC/DEL MC/DEL MC/DEL	8 8 8	ACCOLATE TABS SINGULAIR ² SINGULAIR GRANULES	Use PA Form# 20420 1.Montelukast Granules will only be approved if between ages of 6months-24 months. 2.Singulair Chewables 4mg from 2years-5years and Singulair Chewables 5mgs from 6years-14years old.	
ANTIASTHMATIC - ALPHA-PROTEINASE INHIBITOR				MC MC/DEL MC MC	8 8 8 8	ARALAST ZEMAIRA GLASSIA PROLASTIN SUSR	Use PA Form# 20420	Prolastin and Azemaira will be approved for members with A1AT deficiency and clinically demonstrable panacinar emphysema.
ANTIASTHMATIC - HYDRO-LYTIC ENZYMES				MC/DEL		PULMOZYME SOLN	Use PA Form# 20420	Will be approved for cystic fibrosis patients.
ANTIASTHMATIC - MUCOLYTICS	MC/DEL		ACETYLCYSTEINE ¹	MC		MUCOMYST	1. Acetylcysteine is covered with diagnosis of CF. Use PA Form# 20420	
ANTIASTHMATIC-CFTR POTENTIATOR AND COMBINATIONS				MC MC MC MC MC/DEL		BRONCHITOL ¹ ORKAMBI KALYDECO SYMDEKO TRIKAFTA	1. For the treatment of patients ≥18 years of age with CF. Use PA Form# 20420	<p>Kalydeco will be considered for patients with cystic fibrosis (CF) aged 1 month and older who have at least one mutation in the CFTR gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.</p> <p>Symdeko will be considered for patients with cystic fibrosis (CF) aged 6 years and older who are homozygous for the <i>F508del</i> mutation or who have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor based on in vitro data and/or clinical evidence. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.</p> <p>Bronchitol will be considered as add-on maintenance therapy to improve pulmonary function in adult patients 18 years and older with cystic fibrosis (CF). Use Bronchitol® only for adults</p>

who have passed the Bronchitol® Tolerance Test (BTT). (see Recommended Dosage section for further information)

Trikafta will be considered for the treatment of cystic fibrosis (CF) in patients aged 2 years and older who have at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or mutation in the CFTE gene that is responsive based on in vitro data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to confirm the presence of at least one F508del mutation or a mutation that is responsive based on in vitro data.

Orkambi will be considered for patients with cystic fibrosis (CF) aged 1 year and older who are homozygous for the F508del mutation in the CFTR gene. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of the F508del mutation on both alleles of the CFTR gene. The efficacy and safety of Orkambi have not been established in patients with CF other than those homozygous for the F508del mutation.

[Use PA Form# 20420](#)

IDIOPATHIC PULMONARY FIBROSIS	MC/DEL		OFEV ¹	MC MC		ESBRIET ¹ PIRFENIDONE	1. Diagnosis required Use PA Form# 20420	Ofev- Avoid concomitant use with P-gp and CYP4A inducers (e.g. carbamazepine, phenytoin, and St. John's wort) Esbriet- The concomitant use with strong CYP1A2 inhibitors (e.g. fluvoxamine, enoxacin) is not recommended
COUGH/COLD								
COUGH/COLD	MC/DEL MC/DEL MC/DEL MC MC		DEXTROMETHORPHAN CAPS ¹ DEXTRO-GUAIF SYRP ¹ GUAIFENESIN SYRP ¹ PSEUDOEPHEDRINE ¹ ROBITUSSIN DM SYRP ¹ ROBITUSSIN SUGAR FREE SYRP ¹				1. All of cough cold preparations are not covered except these preferred products. Use PA Form# 20420	All non-preferred products are not covered as permitted by Federal Medicaid regulations and MaineCare Policy.
DIGESTIVE AIDS / ASSORTED GI								
GI - ANTIPERISTALTIC AGENTS	MC/DEL MC/DEL MC/DEL MC		DIPHENOXYLATE DIPHENOXYLATE/ATROPINE LOPERAMIDE HCL CAPS/LIQ OPIUM TINCTURE TINC PAREGORIC TINC	MC/DEL MC MC		LOFENE TABS LONOX TABS MOTOFEN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
GI - ANTI-DIARRHEAL/ ANTACID - MISC.	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ATROPINE SULFATE SOLN BISMATROL BISMUTH SUBSALICYLATE CALCIUM CARBONATE (ANTACID) CHEW DICYCLOMINE HCL GLYCOPYRROLATE TABS HYOSCYAMINE CAPS & TABS HYOSCYAMINE SULFATE KAOPECTATE MAGNESIUM OXIDE TABS MAG-OX 400 TABS PAMINE TABS PROPANTHELINE BROMIDE TABS SODIUM BICARBONATE TABS TUMS	MC/DEL MC/DEL MC MC MC MC MC/DEL MC MC MC MC MC MC MC MC MC MC MC		BELLADONNA ALKALOIDS & OP BENTYL TABS BENTYL SYRP CUVPOSA DARTISLA ODT ² ED-SPAZ MYTESI ¹ GLYCOPYRROLATE INJ LEVSIN TABS LEVSIN/SL SUBL NULEV TBDP OSCIMIN ROBINUL INJ ROBINUL TABS	Use PA Form# 20420 1. Dosing limits apply please refer to Dose Consolidation List 2. It is not indicated as monotherapy for treatment of peptic ulcer because effectiveness in peptic ulcer healing has not been established.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Preferred products that used to require diag codes still require diag codes unless indicated otherwise. Mytesi requires a diagnosis of non-infectious diarrhea in patients with HIV/AIDS on anti-retroviral therapy, prior trials of preferred, more cost effective anti-diarrheals.
GI- BILE ACID				MC		CHOLBAM	Use PA Form# 20420	Indication of bile acid synthesis disorders due to single enzyme defects (SEDs) AND for adjunctive treatment of peroxisomal disorders (PDs)
GI- EOSINOPHILIC ESOPHAGITIS	MC		EOHILIA ¹				Use PA Form# 20420 1. Approvals will not be longer than 12 weeks of treatment in adult and pediatric patients 11 years of age and older	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Eohilia: Dietary modification, PPIs, and topical glucocorticoids are required as initial therapy.
GI - H2-ANTAGONISTS	MC MC/DEL MC/DEL		ACID REDUCER TABS CIMETIDINE FAMOTIDINE	MC MC MC/DEL MC/DEL MC		AXID CAPS AXID AR TABS NIZATIDINE CAPS PEPCID PEPCID AC	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Cimetidine will now be non-preferred and require prior authorization if it is currently being used with any sulfonylurea (except for glyburide). DDI: Cimetidine will require prior authorization if being used in combination with Plavix.

GI - IBAT INHIBITORS				MC MC		BYLVAY ^{1,2} LIVMARLI ^{1,2}	Use PA Form# 20420 1. For the treatment of patients ≥ 3months of age 2. Clinical PA required for appropriate diagnosis	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
GI - PROTON PUMP INHIBITOR	MC/DEL MC/DEL MC/DEL		OMEPRAZOLE CAPS ² PANTOPRAZOLE ² LANSOPRAZOLE CAPS ²	MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	6 6 7 7 8 8 8 8 8 8 8 8 8 8 8	NEXIUM CPDR ³ NEXIUM SUS ⁵ PRILOSEC OTC ³ ACIPHEX TBEC ³ DEXILANT (KAPIDEX) ² KONVOME ² OMEPRAZOLE-SODIUM BICARBONATE CAPS OMEPRAZOLE MAGNESIUM PREVACID CPDR ³ PREVACID SOLUTABS ^{1,4} PRILOSEC CPDR PROTONIX INJ PROTONIX ² VOQUEZNA TABS	1. Prevacid Solutabs available without PA for children less than 9 years old. 2. Dosing limits apply, please see dosage consolidation list. 3. All preferreds and step therapy must be tried and failed. 4. Payment for Prevacid SoluTabs for patients 9 and older will be considered for those patients who cannot tolerate a preferred solid oral dosage form. 5. Nexium sus available without PA if member is < 12 yrs of age and ≤ 1 pack per day Use PA Form# 20720	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to the PPI PA form for additional criteria on Non-Preferred PPIs DDI: Omeprazole will require prior authorization if being used in combination with Plavix. DDI: Lansoprazole will require prior authorization if being used in combination with Plavix. DDI: Prevacid, Omeprazole and pantoprazole will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: Ampicillin, B-12, Fe salts, Griseofulvin, Sporanox, Ketoconazole, Reyataz, or Vantin. DDI: All non-preferred PPIs require prior authorization, but with any prior authorization request, the member's drug profile will also be monitored for current use with ampicillin, B-12, Fe salts, griseofulvin, itraconazole, ketoconazole, Reyataz or Vantin due to a significant drug-drug interaction.
GI - ULCER ANTI-INFECTIVE	MC MC		PYLERA TALICIA			VOQUEZNA DUAL PAK VOQUEZNA TRIPLE PAK	Use PA Form# 20420	
GI - PROSTAGLANDINS	MC		MISOPROSTOL TABS	MC/DEL		CYTOTEC TABS	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
GI - DIGESTIVE ENZYMES	MC/DEL MC		CREON ¹ ZENPEP ¹	MC/DEL MC/DEL MC/DEL		PERTZYE ULTRESA VIOKACE	Use PA Form# 20420 1. Clinical PA is required to establish CF diagnosis and medical necessity. In all cases except cystic fibrosis patients, objective evidence of pancreatic insufficiency (fat malabsorption test etc...) must be supplied.	Non -Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before other non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
GI - ANTI - FLATULENTS / GI STIMULANTS	MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AMITIZA CALULOSE SYRP CONSTULOSE SYRP ENULOSE SYRP GASTROCROM CONC GENERLAC SYRP LACTULOSE SYRP METOCLOPRAMIDE HCL	MC MC/DEL MC MC/DEL		CEPHULAC SYRP INFANTS GAS RELIEF SUSP GIMOTI SPRAY REGLAN TABS		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.

GI - INFLAMMATORY BOWEL AGENTS	MC MC/DEL MC MC MC/DEL MC/DEL	APRISO BALSALAZIDE MESALAMINE ENMA KIT PENTASA SULFAZINE EC TBEC SULFASALAZINE TABS	MC/DEL MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC	ASACOL 800MG HD AZULFIDINE EN-TABS TBEC AZULFIDINE TABS COLAZAL CAPS DELZICOL DIPENTUM CAPS GIAZO LIALDA TABS ¹ MESALAMINE TAB ROWASA ENEM SFROWASA UCERIS RECTAL FOAM ² UCERIS TABS ²	Use PA Form# 20420 Use PA Form# 20420 1. Current users grandfathered. 2. Diagnosis required	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Giazo is only indicated for males, as the safety.efficacy for use in females has not been established.Prior trials of preferred products. Uceris Rectal Foam or Tab- Concomitant use with CYP3A inhibitors (e.g. ketoconazole, itraconazole, ritonavir, indinavir, saquinavir, erythromycin, cyclosporine, and grapefruit juice) should be avoided. Verify prior trials and failures or intolerance of preferred treatments
GI - IRRITABLE BOWEL SYNDROME AGENTS	MC/DEL	LOTROXEN TABS	MC	VIBERZI	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
GI- SHORT BOWL SYNDROME			MC	GATTEX	Use PA Form #20420	Gattex requires a diagnosis of adult SBS who are dependent on parenteral support. Appropriate colonoscopy and lab assessments 6months prior to starting
GI- NASH			MC	REZDIFFRA	Use PA Form #20420	Rezdiffra: The patient must have a diagnosis of NASH with fibrosis Stage 2 or 3 and utilizing imaging and scanning test such as fibro scan, MRI or ultra sound AND the patient does not have evidence of decompensated cirrhosis
MISCELLANEOUS GI						
GI - MISC.	MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL	BISAC-EVAC SUPP BISACODYL BISCOLAX SUPP CINOBAC CAPS CITRATE OF MAGNESIA SOLN CITRUCEL CLENPIQ SOL COLYTE DIOCTO SYRP DOCUSATE CALCIUM CAPS DOCUSATE SODIUM FIBER LAXATIVE TABS FLEET GENFIBER POWD GLYCERIN HIPREX TABS KRISTALOSE PACK LINZESS 145mcg & 290mcg MAALOX MILK OF MAGNESIA SUSP MINERAL OIL OIL MIRALAX BULK POWD (BRAND) MOVANTIK MOVIPREP POWD PACK NULYTELY SOLR PEG 3350- ELECTROLYTE SOL PEG 3350 POWDER SENNA	MC/DEL MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL	ACTIGALL CAPS BENEFIBER CARAFATE CLEARLAX POW COLACE CAPS DIOCTO-C SYRP DOC SOD /CAS CAP DOC-Q-LAX CAPS DOCUSATE SODIUM/CAS CAPS DOK PLUS DULCOLAX SUPP FIBER CON TABS FIBER-LAX TABS GAVILYTE-H GOLYTELY SOLR IBSRELA IQIRVO LINZESS 72mcg ⁴ MALTSUPEX MIRALAX PACKETS MOTTEGRITY OCALIVA ¹ PEG-ELECTROLYTES SOLR PEG 3350 PACKETS PREPOPIK PAK RELISTOR TABS SENXON TABS SENOLOT TABS	1. PA required to confirm FDA approved indication. 2. For the treatment of carcinoid syndrome diarrhea in combination with somatostatin analog (SSA) therapy in adults inadequately controlled by SSA therapy 3. For the treatment of Opioid Induced Constipation(OIC) 4. Established users will be grandfathered	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Linzess is preferred for adults as treatment of IBS-Constipation AND treatment of chronic idiopathic constipation in adults. Trulance should be avoided in pediatric patients less than 18 years of age. Iqirvo: For the treatment of primary biliary cholangitis (PBC) in combination with ursodeoxycholic acid (UDCA) in adults who have had an inadequate response to UDCA, or as monotherapy in patients unable to tolerate UDCA. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

	MC/DEL		SENOKOT GRAN	MC		SENOKOT S TABS		
	MC/DEL		SENOKOT SYRP	MC/DEL		SORBITOL		
	MC/DEL		SENOKOT CHILDRENS SYRP	MC		STOOL SOFTENER PLUS CAPS		
	MC		SENOKOT XTRA TABS	MC		SUFLAVE		
	MC/DEL		STOOL SOFTENER CAPS	MC		SUTAB		
	MC/DEL		SUCRALFATE TABS	MC/DEL		SYMPROIC ³		Use PA Form# 20420
	MC/DEL		SUPREP SOL	MC/DEL		UNI-CENNA TABS		
	MC		TRULANCE ²	MC		UNI-EASE PLUS CAPS		
	MC		UNI-EASE CAPS	MC		V-R NATURAL SENNA LAXATIV TABS		
	MC		URSO FORTE	MC		URSO 250		
	MC/DEL		URSODIOL	MC		XERMELO ²		
MISC. UROLOGICAL								
UROLOGICAL - MISC.	MC		ACETIC ACID 0.25% SOLN	MC		CITRIC ACID/SODIUM CITRAT SOLN	1. Elmiron requires adequate proof of Dx with supportive testing.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC		CYTRA-K SOLN	MC/DEL		CYTRA-2 SOLN		
	MC		FOSFOMYCIN (NDC 82036427401 ONLY)	MC/DEL		ELMIRON CAPS ¹		
	MC		K-PHOS MF TABS	MC		FURADANTIN SUSP		Use PA Form# 20420
	MC/DEL		METHENAMINE MANDELATE TABS	MC/DEL		MACROBID CAPS		
	MC/DEL		NEOSPORIN GU IRRIGANT SOLN	MC/DEL		MACRODANTIN CAPS		
	MC/DEL		NITROFURANTOIN MONO CAPS	MC/DEL		NITROFURANTOIN MACR SUSP		
	MC/DEL		PHENAZOPYRIDINE HCL TABS	MC		POTASSIUM CITRATE/CITRIC SOLN		
	MC/DEL		PHENAZOPYRIDINE PLUS	MC/DEL		PYRIDIUM PLUS TABS		
	MC		POT CITRATE TAB	MC		PYRIDIUM TABS		
	MC/DEL		PROSED/DS TABS	MC/DEL		RENACIDIN SOLN		
	MC		TRICITRATES SYRP	MC		UROCIT-K		
	MC/DEL		URELIEF PLUS					
	MC		UREX TABS					
	MC/DEL		URISED TABS					
	MC/DEL		UROQID #2 TABS					
PHOSPHATE BINDERS								
PHOSPHATE BINDERS	MC/DEL		CALCIUM ACETATE CAP ¹	MC		AURYXIA ¹	Use PA Form# 20420	Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before less preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		FOSRENOL CHEW ¹	MC/DEL		CALCIUM ACETATE TAB ¹	1. Diag required.	
	MC/DEL		MAGNEBIND - 400 ¹	MC/DEL		ELIPHOS ¹		
	MC		PHOSLYRA ¹	MC/DEL		FOSRENOL PWDR ¹		
	MC/DEL		REVELA ¹	MC		VELPHORO ¹		Xphozah to reduce serum phosphorus in adults with chronic kidney disease (CKD) on dialysis as add-on therapy in patients who have an inadequate response to phosphate binders or who are intolerant of any dose of phosphate binder therapy.
	MC/DEL			MC		XPHOZAH		
INTRA-VAGINALS								
VAGINAL - ANTIBACTERIALS	MC/DEL		CLEOCIN CREA	MC/DEL		METROGEL VAGINAL GEL ¹	1. Dosing limits apply, please see Dosage Consolidation List.	Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before less preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		CLEOCIN SUPP	MC/DEL		VANAZOLE		
	MC		CLINDESSE CREA	MC		XACIATO		
	MC/DEL		METRONIDAZOLE VAGINAL GEL ¹					
	MC/DEL		NUVESSA				Use PA Form# 20420	
VAGINAL - ANTI FUNGALS	MC/DEL		CLOTRIMAZOLE CREA	MC		AVC CREA	1. Quantity limit: 1/script/2 weeks	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		CLOTRIMAZOLE-3 CREA	MC		CLOTRIMAZOLE 3 DAY CREA		
	MC/DEL		GYNE-LOTRIMIN CREA	MC		GYNAZOLE-1 CREA		
	MC		MICONAZOLE CREA	MC		GYNE-LOTRIMIN 3 TABS		
	MC		MICONAZOLE 3 KIT CREA OTC	MC/DEL		MICONAZOLE 3 COMBO PACK KIT ¹		
	MC/DEL		MICONAZOLE 7 CREA	MC/DEL		MICONAZOLE 3 SUPP		DDI: Miconazole will require prior authorization if being used in combination with Warfarin.
	MC/DEL		MICONAZOLE NITRATE CREA	MC		TERAZOL 3 CREA		
	MC		NYSTATIN TABS	MC		TERAZOL 7 CREA		
	MC/DEL		TERCONAZOLE CREAM	MC/DEL		TERCONAZOLE SUPP		
	MC		VAGITROL					

	MC		V-R MICONAZOLE-7 CREA						
VAGINAL - CONTRACEPTIVES									Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Use PA Form# 20420
VAGINAL - ESTROGENS	MC/DEL MC/DEL		ESTRING RING PREMARIN CREA	MC/DEL MC/DEL		ESTRACE CREA ¹ VAGIFEM TABS ¹	1. Must fail all preferred products before non-preferred. Use PA Form# 20420		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
VAGINAL - OTHER	MC/DEL MC MC		ACID JELLY GEL ACI-JEL GEL CERVICAL AMINO ACID CREA	MC		AMINO ACID CERVICAL CREA	Use PA Form# 20420		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BENIGN PROSTATIC HYPERPLASIA (BPH)									
BPH	MC/DEL MC/DEL MC/DEL MC/DEL		DOXAZOSIN MESYLATE TABS FINASTERIDE ¹ 5mg TERAZOSIN HCL CAPS TAMSULOSIN HCL	MC/DEL MC/DEL MC MC MC/DEL MC/DEL	5 8 8 8 8 8	FLOMAX CP24 ALFUZOSIN AVODART ^{2,4} CARDURA TABS ⁴ ENTADFI ^{5,6} JALYN ^{3,4} PROSCAR TABS ⁴ RAPAFLO ⁴ UROXATRAL ⁴	1. There will be dosing limits of 1 tab per day with out PA. 2. Prior use of preferred agent prior to any approvals. 3. Use of preferred (tamsulosin and finasteride) and (tamsulosin and non-preferred Avodart). 4. Non-preferred products must be used in specified order. 5. Use of individual ingredients preferred (Finasteride and tadalafil). 6. Entadfi® is not recommended for more than 26 weeks Use PA Form# 20420		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approval of a non-preferred 5-alpha reductase inhibitor requires objective clinical evidence of a very enlarged prostate rather than just the presence of obstructive urinary outflow symptoms along with adequate trial of preferred Proscar.
ANXIOLYTICS									
ANXIOLYTICS - BENZODIAZEPINES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ALPRAZOLAM TABS CHLORDIAZEPOXIDE HCL CAPS CLORAZEPATE DIPOTASSIUM TABS DIAZEPAM LORAZEPAM OXAZEPAM CAPS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL	8 8 8 8 8 8 9	ALPRAZOLAM ER ATIVAN LOREEV XR NIRAVAM SERAX TRANXENE XANAX TABS XANAX XR	Use PA Form# 20420		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANXIOLYTICS - MISC.	MC/DEL MC MC MC/DEL MC/DEL MC/DEL		BUSPIRONE HCL TABS HYDROXYZINE HCL SOLN HYDROXYZINE HCL SYRP HYDROXYZINE HCL TABS ¹ HYDROXYZINE PAMOATE CAPS MEPROBAMATE TABS	MC MC MC/DEL MC/DEL MC/DEL		BUSPAR TABS DROPERIDOL SOLN DROPERIDOL SOLN DROPERIDOL SOLN	Use PA Form# 20420 1. Dosing limits apply, please refer to Dose consolidation list.		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTI-DEPRESSANTS									
ANTIDEPRESSANTS - MAO INHIBITORS	MC/DEL		NARDIL TABS	MC/DEL		TRANLYCYPROMIINE	Use PA Form# 20420		

ANTIDEPRESSANTS - MAO INHIBITORS TOPICAL				MC/DEL		EMSAM ¹	1. Dosing limits apply, please refer to Dose consolidation list. Use PA Form# 20420	Preferred drugs (including a preferred SSRI, a non-SSRI, and Venlafaxine ER) must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIDEPRESSANTS - SELECTED SSRI's AND OTHERS	MC/DEL		BUPROPION HCL TABS	MC/DEL	8	APLENZIN ⁴	1. Strong caution with pediatric population.	Preferred drugs (including failure of at least one preferred SSRI, one SNRI and one non-SSRI/SNRI) must be tried for at least 4 weeks each and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		BUPROPION SR	MC	8	AUVELITY ¹¹		
	MC/DEL		BUPROPION XL 150mg and 300mg	MC/DEL	8	BUPROPION XL 450mg	2. Max daily dose allowed is 120mg. Combination of multiple strengths require	
	MC/DEL		CITALOPRAM	MC/DEL	8	CELEXA	PA	
	MC/DEL		DULOXETINE ^{2,9}	MC	8	CYMBALTA ²	4. Dosing limits allowing 2 tabs/day and a max daily limit of 200mg / day applies. Please see dose consolidation list.	CYMBALTA: Fibromyalgia diagnosis- prior use and failure of preferred generics (amitriptyline or cyclobenzaprine) <u>and</u> gabapentin prior to approval.
	MC/DEL		ESCITALOPRAM	MC/DEL	8	DRIZALMA SPRINKLES		
	MC/DEL		FLUOXETINE 10mg AND 20mg AND 40mg CAPS	MC/DEL	8	EFFEXOR TABS		
	MC/DEL		FLUOXETINE HCL LIQD	MC/DEL	8	EFFEXOR XR CP24		
	MC/DEL		FLUVOXAMINE MALEATE TABS	MC/DEL	8	FETZIMA ⁷		
	MC/DEL		MIRTAZAPINE	MC/DEL	8	FLUOXETINE 10mg AND 20mg AND 60mg TABS	5. Dosing limits apply, please refer to Dose consolidation list and max daily dose applies. Max daily dose allowed is 375mg.	DDI: Fluvoxamine will now be non-preferred and require prior authorization if it is currently being used with glimepiride (Amaryl).
	MC/DEL		NEFAZODONE	MC	8	FORFIVO XL		DDI: Preferred nefazodone will now be non-preferred and require prior authorization if it is currently being used in combination with either Onglyza 5mg, Enablex 15mg or Vesicare 10mg.
	MC/DEL		PAROXETINE ¹	MC/DEL	8	IRENKA		
	MC/DEL		SERTRALINE HCL	MC/DEL	8	KHEDEZLA		
	MC/DEL		TRAZODONE HCL TABS	MC/DEL	8	LEXAPRO TABS		
	MC/DEL		VENLAFAXINE ER CAPS ⁵	MC	8	LUVOX TABS	6. Non-preferred products must be used in specified step order.	DDI: Fluoxetine will require prior authorization if being used in combination with Plavix. DDI: Fluvoxamine will require prior authorization if being used in combination with Plavix.
	MC/DEL		VENLAFAXINE TABS ⁵	MC	8	MAPROTILINE HCL TABS		
				MC/DEL	8	MIRTAZAPINE ODT		
				MC	8	OLEPTRO		
				MC/DEL	8	PAROXETINE CR ¹	7. Requires previous trials/failure of multiple preferred medications.	SAVELLA: Fibromyalgia diagnosis and trial of a preferred generic amitriptyline, cyclobenzaprine, duloxetine and gabapentin prior to approval.
				MC/DEL	8	PAXIL ¹	Dosing limits apply, please see the dose consolidation list. Max daily dose of 80mg if used concomitantly with strong CYP3A4 inhibitor.	DDI: Drizalma Sprinkle avoid the concomitant use of duloxetine with potent CYP1A2 inhibitors (e.g. fluvoxamine, cimetidine, ciprofloxacin, enoxacin).
				MC/DEL	8	PAXIL CR ¹		
				MC	8	PRISTIQ		Zulresso® is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the Zulresso® REMS.
				MC	8	PROZAC		
				MC	8	PROZAC CAPS		
				MC	8	PROZAC WEEKLY CPDR		
				MC/DEL	8	REMERON TABS	8. Psychiatry recommended. Please see criteria section.	Spravato: Treatment Resistant Depression
				MC/DEL	8	SARAFEM CAPS		• Must be 18 years of age or older; and medication must be administered under the direct, on site, supervision of a licensed healthcare provider with post-administration observation of a minimum of least 2-hours. The medication must be prescribed by or in consultation with a psychiatrist and prescriber must be enrolled in the REMS program.
				MC/DEL	8	SPRAVATO ⁸		• Approval is based upon failure of at least two antidepressants and failure of an antidepressant used adjunctively with one recognized augmentation strategy such as lithium, an atypical antipsychotic, thyroid hormone, etc
				MC/DEL	8	TRAZODONE HCL 300MG TABS	9. Please use multiples of the 20mg, the 40mg is still non-preferred.	• Ongoing use of Spravato beyond 3 months is based upon a positive response as evidenced by at least a 30 % reduction from baseline as measured by a standardized rating scale
				MC/DEL	8	TRINTELLIX		
				MC	8	WELLBUTRIN TABS	10. For the treatment of patients ≥ 18 years of age.	
				MC	8	WELLBUTRIN SR TBCR		
				MC	8	WELLBUTRIN XL		
				MC/DEL	8	REMERON SOLTAB TBDP	11. Use individual ingredients separately.	Spravato: MDD with Suicidal Ideation Approval for this indication only if it is started in an inpatient unit, given adjunctively with an optimized antidepressant regimen, and with an 8-12 week initial approval with ongoing use dependent upon documentation of ongoing benefit.
				MC/DEL	8	SAVELLA ⁴		
				MC/DEL	8	ZOLOFT	12. Approval will be limited to a 14-day treatment course.	DDI: Reduce the Zurzuvae® dosage when used with a strong CYP3A4 inhibitor.
				MC/DEL	8	ZULRESSO ¹⁰		
				MC	8	ZURZUVAE ¹²		
				MC/DEL	8	VENLAFAXINE ER TABS ⁵		
				MC/DEL	9	VIIBRYD ⁵		
				MC/DEL	9	FLUOXETINE 90mg TABS ⁵	Use PA Form# 20420	
ANTIDEPRESSANTS - TRI-CYCLICS	MC/DEL		AMITRIPTYLINE HCL TABS ¹	MC/DEL		AMOXAPINE TABS	1. Users over the age of 65 require a pa.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		CLOMIPRAMINE HCL CAPS ¹	MC/DEL		ANAFRANIL CAPS		
	MC/DEL		DESIPRAMINE HCL TABS ¹	MC/DEL		DOXEPIN HCL 150 MG ²		
	MC/DEL		DOXEPIN HCL ¹ (not generic Silenor)	MC/DEL		DOXEPIN (generic Silenor)		
	MC/DEL		IMIPRAMINE HCL TABS ¹	MC/DEL		NORPRAMIN TABS	2. Use multiples of 50mg.	
	MC/DEL		NORTRIPTYLINE HCL ¹	MC/DEL		PAMELOR		
	MC		PROTRIPTYLINE HCL TABS ¹	MC		TOFRANIL	Use PA Form# 20420	
	MC		SURMONTIL CAPS ¹	MC		VIVACTIL TABS	Use PA Form# 10220 for Brand Name requests	

SEDATIVE / HYPNOTICS								
SEDATIVE/HYPNOTICS - BARBITURATE	MC MC/DEL MC MC/DEL		BUTISOL SODIUM TABS ¹ CHLORAL HYDRATE SYRP ¹ MEBARAL TABS ¹ PHENOBARBITAL ¹	MC MC/DEL		LUMINAL SOLN SOMNOTE CAPS	1. PA required for new users of preferred products if over 65 years. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
SEDATIVE/HYPNOTICS - BENZODIAZEPINES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		DORAL TABS ¹ ESTAZOLAM TABS ¹ FLURAZEPAM HCL CAPS ¹ TEMAZEPAM CAPS 15 & 30MG ¹ TRIAZOLAM TABS ¹	MC MC MC/DEL MC/DEL		HALCION TABS ¹ MIDAZOLAM HCL SYRP RESTORIL CAPS ¹ TEMAZEPAM 7.5MG ¹	1. Dosing limits apply, please see dosing consolidation list. Use PA Form# 30110	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Benzodiazepines do cause dependence with continued use and usage should be limited to 7-10 days at a time. Chronic intermittent use (2-3 Days per week max) is the standard of care
SEDATIVE/HYPNOTICS - Non-Benzodiazepines	MC/DEL MC MC/DEL MC/DEL	1 1 1 2	MIRTAZAPINE TRAZODONE ZOLPIDEM ² ZALEPLON ^{2,3}	MC/DEL MC/DEL MC/DEL MC/DEL	7 7 7 8 8 8 8 8 8 8 8 8	AMBIEN ¹ ESZOPICLONE ZOLPIDEM ER AMBIEN CR ¹ BELSOMRA ¹ DAYVIGO ¹ EDLUAR HETLIOZ INTERMEZZO LUNESTA ¹ SONATA CAPS ¹ ROZEREM QUVIVIQ ZOLPIMIST	1. Quantity Limit of 12 per 34 days. 2. Quantity limits will be allowed up to 30/30, but intermittent therapy is recommended. 3. Only zolpidem trial/failure will be required to obtain Zaleplon. 4. Must fail all preferred products before non-preferred Use PA Form# 30110	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Ambien, Ambien CR, Lunesta, Sonata, Zaleplon and Zolpidem may cause dependence with continued use and as with benzodiazepines, usage should be limited to 7-10 days at a time. Chronic intermittent use (2-3 days per week max) is the standard of care. Please refer to Sedative/Hypnotic PA form. DDI: Belsonra® with strong CYP3A inhibitors (e.g. ketoconazole, itraconazole, posaconazole, clarithromycin, nefazodone, ritonavir, saquinavir, nelfinavir, indinavir, boceprevir, telaprevir, telithromycin, and conivaptan) is not recommended
ANTI-PSYCHOTICS								
ANTIPSYCHOTICS - ATYPICALS	MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		ABILIFY MAINTENA ARIPIPRAZOLE TAB ³ ARISTADA ARISTADA INITIO OLANZAPINE ^{2,3} OLANZAPINE ^{2,3} ODT INVEGA HAFYERA INVEGA SUSTENNA INVEGA TRINZA INJ LURASIDONE TAB PALIPERIDONE ER PERSERIS RISPERDAL CONSTA RISPERIDONE ODT RISPERIDONE TAB ^{2,3} RISPERIDONE SOLN ² QUETIAPINE ^{2,3} QUETIAPINE XR VRAYLAR ⁴ ZIPRASIDONE ^{2,3}	MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC	8 8	ABILIFY ASIMTUFII ABILIFY DISC TAB, INJ and SOL ¹ ABILIFY TABS ² ARIPIPRAZOLE SOL ARIPIPRAZOLE ODT CAPLYTA FANAPT GEODON INVEGA IGALMI LATUDA LYBALVI NUPLAZID REXULTI RISPERDAL TAB RISPERDAL M TAB ¹ RISPERDAL SOLN RYKINDO SAPHRIS ¹ SECUADO SEROQUEL TABS UZEDY ZYPREXA TABS	If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine. This also includes combination of Seroquel with Seroquel XR. Use PA form# 20440 for Multiple Antipsychotic requests Use PA form# 10130 for non preferred single therapy atypical requests 1. Established users of single therapy atypicals were grandfathered. 2. Prior Authorization will be required for preferred medications for members under the age of 5	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Non preferred atypicals will be approved for patients with FDA-approved indications, and for specific conditions supported by at least two published peer-reviewed double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality and as long as all first line preferred therapies have been tried and failed at full therapeutic doses for adequate durations (at least two weeks). Prescriptions for quetiapine are limited to a maximum daily dose of 800mg. Uzedy: Establish tolerability with oral risperidone prior to initiating Uzedy Atypicals: Prior Authorization will be required for preferred medication to assure indication is in accordance with FDA approved or literature supported evidence-based best practices. The approved indications are: <ul style="list-style-type: none"> schizophrenia bipolar disorder agitation related to autism adjunct in major depressive disorder If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine. This also includes combination of Seroquel with Seroquel XR. DDI: It is recommended to reduce the Vraylar® dose if it is used concomitantly with a strong CYP3A inhibitor (such as itraconazole, ketoconazole). The concomitant use of Vraylar® with a CYP3A4 inducer (such as rifampin, carbamazepine) is not recommended.

				MC MC MC/DEL	8 9	ZYPREXA RELPREVV ZYPREXA ZYDIS TBDP ¹ SEROQUEL XR		3. Dosing limits apply please refer to the dose consolidation list. 4.Requires step through 1 preferred drug for all indications except AMDD. AMDD requires insufficient response from two antidepressants	DDI: The concomitant use of Nuplazid with other drugs known to prolong the QT interval (e.g. Class IA antiarrhythmics, Class 3 antiarrhythmics, antipsychotics, and antibiotics such as gatifloxacin and moxifloxacin). Lybalvi: Step through aripiprazole and Latuda. If criteria is met then initial approval for 3 months. Subsequent approvals will be based on evidence of not gaining >= 10 % baseline body weight for ongoing approval. If weight gain >= 10 % of initial body weight, then criteria for ongoing use not met. Invega Hafyera: The patient is started and stabilized on the medication OR The patient has been adequately treated with Invega Sustenna (paliperidone palmitate 1-month) for at least four months or Invega Trinza (paliperidone palmitate 3- month) following at least one 3-month injection cycle.	
ANTIPSYCHOTICS - SPECIAL ATYPICALS	MC/DEL		CLOZAPINE TABS	MC/DEL MC/DEL MC/DEL		CLOZAPINE ODT CLOZARIL TABS VERSACLOZ SUSP		Use PA Form# 20420	Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred brand will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Patients previously stabilized on brand name drug will be approved.	
ANTIPSYCHOTICS - TYPICAL	MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		CHLORPROMAZINE HCL FLUPHENAZINE DECANOATE FLUPHENAZINE HCL HALDOL HALOPERIDOL HALOPERIDOL DECANOATE SOLN HALOPERIDOL LACTATE SOLN LOXAPINE SUCCINATE CAPS LOXITANE-C CONC MOBAN TABS PERPHENAZINE PROCHLORPERAZINE SERENTIL THIORIDAZINE HCL THIOTHIXENE TRIFLUOPERAZINE HCL TABS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		COMPAZINE COMPRO SUPP FLUPHENAZINE HCL CONC HALDOL DECANOATE LOXITANE CAPS MELLARIL NAVANE CAPS PROLIXIN STELAZINE TABS		Use PA Form# 20420 If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine.	
LITHIUM										
LITHIUM	MC/DEL MC/DEL		LITHIUM CARBONATE LITHIUM CITRATE SYRP	MC/DEL MC/DEL		ESKALITH CAPS ESKALITH CR TBCR		Use PA Form# 20420		
COMBINATION - PSYCHOTHERAPEUTIC										
PSYCHOTHERPEUTIC COMBINATION	MC/DEL MC/DEL		CHLORDIAZEPOXIDE/AMITRIPT PERPHENAZINE/AMITRIPTYLIN					Use PA Form# 20420		
STIMULANTS										
STIMULANT - AMPHETAMINES -SHORT ACTING	MC/DEL MC/DEL MC		AMPHETAMINE SALT COMBO ^{1,4} DEXTROAMPHET SULF TABS PROCENTRA	MC/DEL MC MC/DEL MC		ADDERALL TABS EVEKEO METHAMPHETAMINE HCL ZENZEDI		1. Preferred stimulants will be available without PA if diagnosis of ADHD or Narcolepsy.		

						<p>2. As per recent FDA alert, Adderal & Dexedrine should not be used in patients with underlying heart defects since they may be at increased risk for sudden death.</p> <p>3. Dosing limits apply, please see dosing consolidation list.</p> <p>4. Max daily dose of 50mg.</p> <p>Use PA Form# 20420</p>	
STIMULANT - LONG ACTING AMPHETAMINES SALT	MC/DEL MC MC		<p>AMPHETAMINE/DEXTROAMPHET ER^{3,4,7}</p> <p>ADDERALL XR CP24^{1,3,4,7}</p> <p>VYVANSE^{2,3,4}</p>	MC MC MC		<p>MYDAYIS⁵</p> <p>VYVANSE CHEW^{2,3,4,6}</p> <p>XELSTRYM⁸</p> <p>Use PA Form# 20420</p> <p>1. As per recent FDA alert, Adderall should not be used in patients with underlying heart defects since they may be at increased risk for sudden death.</p> <p>2. FDA approval is currently for adults and children 6 or older. Will be available without PA for this age group if within dosing limits. Limit of one capsule daily. Max dose of 70MG daily.</p> <p>3. Preferred stimulants will be available without PA if diagnosis of ADHD.</p> <p>4. Dosing limits apply, please see dosing consolidation list.</p> <p>5. For the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in patients 13 years and older</p> <p>6. Vyvanse chew grace period for current user through June 2022.</p> <p>7. FDA approval is currently for adults and children 6 or older. Will be available without PA for this age group if within dosing limits. Max dose of 50MG daily without a PA.</p> <p>8. For the treatment of patients 6 years of age and older.</p>	<p>DDI: The concomitant use of Mydayis® is contraindicated with monoamine oxidase inhibitors (MAOIs) or within 14 days after discontinuing MAOI treatment, as concomitant use can increase hypertensive crisis.</p>
LONG ACTING AMPHETAMINES	MC MC/DEL		<p>DEXTROAMPHET SULF CPSR^{1,3}</p> <p>DEXTROAMPHETAMINE ER</p>	MC/DEL MC		<p>ADZENYS ER³</p> <p>ADZENYS XR- ODT</p> <p>1. Preferred stimulants will be available without PA if diagnosis of ADHD.</p>	

	MC	DYANAVEL XR SUS	MC MC MC	ADZENYS XR ³ DEXEDRINE CAP SR ^{2,3} DYANAVEL XR TAB	2. As per recent FDA alert, Adderall & Dexedrine should not be used in patients with underlying heart defects since they may be at increased risk for sudden death. 3. Dosing limits apply, please see dosing consolidation list. Use PA Form# 20420	DDI: : The concomitant use of Adzenys® XR is contraindicated with monoamine oxidase inhibitors (MAOIs) or within 14 days after discontinuing MAOI treatment.
STIMULANT - METHYLPHENIDATE	MC/DEL MC/DEL MC/DEL MC/DEL	DEXMETHYLPHENIDATE IR TABS METHYLPHENIDATE SOL METHYLPHENIDATE TAB METHYLIN TABS ^{1,2}	MC/DEL MC/DEL MC MC MC/DEL MC/DEL	FOCALIN IR TABS METADATE ER METHYLPHENIDATE HCL CHEW METHYLIN CHEWABLES METHYLIN SOL RITALIN	1. Preferred stimulants will be available without PA if diagnosis of ADHD. Use PA Form# 20420 2. Dosing limits apply, please see dosing consolidation list. Maximum daily doses are as follows: 72mg daily for methylphenidate and 36mg daily for dexamethylphenidate.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E.
STIMULANT - METHYLPHENIDATE - LONG ACTING	MC MC/DEL MC MC MC/DEL	CONCERTA TBCR DEXMETHYLPHENIDATE CAP ER 50/50 QUILLICHEW ER ^{5,1} QUILLIVANT XR SUS ^{1,5} RITALIN LA ⁴	MC MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	5 METADATE CD CPR 8 ADHANSIA XR ^{2,6} 8 APTENSIO XR ² 8 AZSTARYS ⁶ 8 COTEMPLA XR ² 8 COTEMPLA XR ODT ² 8 DAYTRANA ^{2,3} 8 FOCALIN XR ² 8 JORNAY PM ^{2,6} 8 METHYLPHENIDATE ER CAPS ^{2,4} 8 METHYLPHENIDATE LA CAPS ² 8 METHYLPHENIDATE ER ^{2,4} CAPS 50/50 8 METHYLPHENIDATE ER ² CAPS 40/60 8 METHYLPHENIDATE CD CAPS ² 30-70	1. Preferred stimulants will be available without PA if diagnosis of ADHD. 2. Non-preferred products must be used in specified step order. 3. FDA approval currently only for ages 6-16. Limit of one patch daily. Max dose of 30MG daily. 4. Dosing limits apply, please see dosing consolidation list 5. Quillivant XR and Quillichew ER are only indicated for use in patients 6 years of age and older. 6. For the treatment of patients ≥ 6 years of age. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
STIMULANT - STIMULANT LIKE	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	ATOMOXETINE HCL ARMODAFINIL CLONIDINE ER GUANFACINE ER MODAFINIL TABS QELBREE ^{6,7}	MC/DEL MC MC MC/DEL MC MC/DEL MC	7 PROVIGIL TABS ³ 7 STRATTERA ^{1,2} 8 CAFKIT SOLN ³ 8 INTUNIV 8 KAPVAY 8 SUNOSI 8 WAKIX	1. Failure of both an amphetamine and methylphenidate is required for consideration for approval of Strattera, unless history of substance abuse without current use of abusable medication(s). Additionally, for patients <17 years of age, a trial of	Provigil requests require diagnosis of Narcolepsy, ADHD, or Obstructive Sleep Apnea. Previous failures of methylphenidate and amphetamine is required for Narcolepsy and ADHD diagnosis, with additional Strattera trial needed with ADHD diagnosis. Please refer to detailed criteria on Provigil PA form Sunsosi is non-preferred and is indicated for to improve wakefulness in adult patients with excessive daytime sleepiness associated with narcolepsy or obstructive sleep apnea (OSA). Wakix is non-preferred and is indicated for the treatment of excessive daytime sleepiness (EDS) in adults with narcolepsy DDI: Sunosi® is contraindicated with MAO inhibitors or within 14 days after discontinuing the MAO inhibitor.

				MC	8	XYREM SOL	<p>quanfacine in required before approval of Strattera.</p> <p>2. Strattera currently has dosing limitations allowing one tablet per day for all strengths if obtain approval. Max daily dose of Strattera is 100mg. Please see dosing consolidation list.</p>	
				MC	8	XYWAV ⁵		<p>Xywav: Diagnosis of cataplexy associated with narcolepsy OR excessive daytime sleepiness associated with narcolepsy. Diagnosis must be confirmed by submission of supporting documentation to include the specialist's interpretation of the Polysomnography (PSG) and Multiple Sleep Latency Test (MSLT) results</p> <p>FDA reminded healthcare professionals and patients that the combined use of Xyrem (sodium oxybate) with alcohol or central nervous system (CNS) depressant drugs can markedly impair consciousness and may lead to severe breathing problems (respiratory depression)</p> <p>DDI: Concomitant use of Qelbree® with an MAO inhibitor or within 2 weeks after discontinuing an MAO inhibitor is contraindicated</p> <p>DDI: Concomitant use of Qelbree® significantly increases the total exposure, but not peak exposure, of sensitive CYP1A2 substrates, which may increase the risk of adverse reactions associated with these CYP1A2 substrates. Coadministration of Qelbree® with sensitive CYP1A2 substrates or CYP1A2 substrates with a narrow therapeutic range (e.g. alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, theophylline), is contraindicated.</p> <p>Use PA Form# 20710 for Provigil, Nuvigil and Xyrem</p> <p>Use PA Form# 20420 for all others</p>
				MC/DEL	9	NUVIGIL ³	3. Non-preferred products must be used in specified	
				MC	9	DESOXYN TABS ³	4. Please use generic Guanfacine.	
				MC	9	DESOXYN CR ³	5. For patients 7 years of age and older with 6. For pediatric patients 6 years of age or older 7. Preferred with a trial and fail either Atomoxetine OR any 2 preferred ADHD agents.	

ANTI-CATAPLECTIC AGENTS

PSYCHOTHERAPEUTIC AGENTS - MISC.				MC		NUDEXTA		
				MC		XENAZINE		Use PA Form# 20710 for Xenazine

WEIGHT LOSS

WEIGHT LOSS							No longer covered: PHENTERMINE, XENICAL, DIDREX, and MERIDIA	Weight loss drugs are not covered as permitted by Federal Medicaid regulations and Maine Medicaid (MaineCare) Policy.
-------------	--	--	--	--	--	--	--	---

ALZHEIMER DISEASE

ALZHEIMER - Cholinomimetics/Others	MC/DEL		DONEPEZIL HYDROCHLORIDE TABS ¹	MC	6	ARICEPT TABS ²	<p>1. PA is required to establish dementia diagnosis and baseline mental status score.</p> <p>2. Must fail all preferred products before moving to non-preferred.</p> <p>3. Approvals will require trials and failure or clinical rationale why preferred patches cant be used.</p>	<p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Kisunla and Leqembi: Testing to rule out reversible causes of dementia (CBC, CMP, TSH, B12, urine drug screen, RPR/VDRL, (folate (if alcohol abuse is present), HIV (if risk present) and an assessment including a review of current medications as a cause of intellectual decline</p> <p>- Prescribed by or in consultation with a neurologist or geriatrician or geriatric psychiatrist. Diagnosis of Alzheimer's disease defined as:</p> <ul style="list-style-type: none"> •Confirmed presence of amyloid pathology and mild cognitive impairment or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease OR •Confirmed presence of amyloid pathology and prodromal or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease <p>-Testing:</p> <ul style="list-style-type: none"> •Clinical Dementia Rating (CDR) global score of 0.5 or 1.0 OR •Repeatable Battery for Assessment of Neuropsychological Status (RBANS) delayed memory index score ≤ 85 OR •Mini-Mental State Examination (MMSE) score of 20-30 OR •Montreal Cognitive Assessment (MoCA) score ≤ 22 <p>- Member is age 50 or older</p> <p>- Obtain recent (within one year) brain magnetic resonance imaging (MRI) prior to initiating treatment</p> <p>- Provider attestation to obtain MRIs prior to the 7th infusion (first dose of 10 mg/kg) and 12th infusion (sixth dose of 10 mg/kg)</p>
	MC/DEL		DONEPEZIL HYDROCHLORIDE ODT ¹	MC	6	ARICEPT ODT ²		
	MC/DEL		EXELON DIS ¹	MC/DEL	7	DONEPEZIL HYDROCHLORIDE TABS 23MG		
	MC/DEL		GALANTAMINE CAPS ¹	MC	8	ADLARITY ³		
	MC/DEL		GALANTAMINE TAB ¹	MC/DEL	8	EXELON CAP		
	MC/DEL		MEMANTINE ¹	MC/DEL	8	GALANTAMINE HYDROBROMIDE SOL		
	MC/DEL		RIVASTIGMINE TARTRATE CAPS ¹	MC	8	KISUNLA		
				MC	8	LEQEMBI ^{1,2}		
				MC/DEL	8	MEMANTINE HCL SOL		
				MC/DEL	8	NAMENDA		
				MC/DEL	8	NAMENDA XR CAPS		
				MC/DEL	8	NAMZARIC		
				MC	8	RAZADYNE ²		

				MC	9	COGNEX CAPS ²		<p>Member does NOT have history or increased risk of amyloid related imaging abnormalities-edema (ARIA-E), which includes brain edema or sulcal effusions and amyloid related imaging abnormalities hemosiderin deposition (ARIA-H), which includes microhemorrhage and superficial siderosis</p> <p>Member does NOT have hypersensitivity to any components of these drugs</p> <p>Failure of or inability to tolerate at least two other preferred Alzheimer therapies for at least four months each, one of which should include a combination of a cholinesterase inhibitor with memantine</p> <p>If the initial drug utilized is the combination of a cholinesterase inhibitor and memantine, then only that single trial of two drugs is required</p>
SMOKING CESSATION								
NICOTINE PATCHES / TABLETS	MC/DEL MC/DEL MC/DEL MC/DEL		CHANTIX TAB ¹ CHANTIX STARTER PACK NICOTINE DIS PT24 ¹ VARENICLINE TAB	MC/DEL		NICODERM CQ PT24 ¹	Use PA Form# 20420 1. See criteria section for exemptions	<p>As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines.</p> <p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations</p> <p>Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.</p>
NICOTINE REPLACEMENT - OTHER	MC/DEL MC/DEL MC/DEL		NICOTINE POLACRILEX GUM ¹ NICOTINE LOZENGE MINI NICOTINE LOZENGE	MC/DEL MC/DEL MC/DEL MC	8 8 8 8	NICOTROL INHALER ^{1,2} NICOTROL NASAL SPRAY ^{1,2} NICORETTE GUM ^{1,2} NICORETTE LOZENGES	Use PA Form# 20420 1. See criteria section for exemptions 2. Must use non-preferred products in specified step order.	<p>As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines.</p> <p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations</p> <p>Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.</p>
ALCOHOL DETERRENTS								
ALCOHOL DETERRENTS	MC/DEL MC MC MC/DEL		ACAMPROSATE ANTABUSE TABS DISULFIRAM TABS NALTREXONE HCL TABS	MC/DEL		ACAMPRO ¹	1. Should only be used in conjunction with formal structured outpatient detoxification program. Use PA Form# 20420	Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS ANALGESICS								
ANALGESICS - MISC.	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		ACETAMINOPHEN ASPIRIN ASPRIN/ APAP/ CAFF TAB BUTAL/ASA/CAFF BUTALBITAL COMPOUND BUTALBITAL/ACET TABS BUTALBITAL/APAP CAPS BUTALBITAL/APAP/CAFFEINE TABS CHOLINE MAGNESIUM TRISALI DIFLUNISAL TABS EXCEDRIN SALSALATE TABS	MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC MC		AXOCET CAPS ESGIC-PLUS FIORICET TABS FIORINAL CAPS FIORTAL CAPS FORTABS TABS PHRENILIN TABS PHRENILIN FORTE CAPS TRILISATE LIQD TRILISATE TABS ZEBUTAL CAPS ZORPRIN TBCR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
LONG ACTING NARCOTICS								
NARCOTICS - LONG ACTING	MC/DEL MC/DEL		FENTANYL PATCH ⁴ BUTRANS ⁴	MC MC	8 8	ARYMO ER AVINZA	Use PA Form# 20510 Use PA form #10300 for	Preferred drugs (Fentanyl Patch, Morphine Sulfate ER tab, Butrans and Embeda) must be tried for at least 2 weeks each & failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Adequate clinical dose/ratio/treatment of non-preferred drug

	MC/DEL MC/DEL MC	MORPHINE SULFATE ER TB12 NUCYNTA ER XTAMPZA ER	MC MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC MC MC/DEL	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 9	BELBUCA EXALGO HYSINGLA ER KADIAN METHADONE METHADOSE MORPHABOND ER MORPHINE SULFATE ER CAP MORPHINE SULFATE SUPP MS CONTIN TB12 OPANA ER ORAMORPH SR TB12 OXYCONTIN TB12 ¹ XARTEMIS ER ZOHYDRO ER OXYCODONECONC OXYCODONE ER ^{3,5}	PAs over the opiate limit 1. Oxycontin will be available without PA for patients treated for or dying from cancer or hospice patients. CA (cancer) or HO (hospice) diag code may be used but store must verify since all scripts will be audited and stores will be liable. 2. Established users are grandfathered. 3. Oxycodone ER allowed only 2 per day for all strengths except 80 mg, please see attached to 4. Dosing limits apply. Please see dose consolidation list. 5. Non-preferred products must be used in specific order. 6. Methadone will be available without PA for patients treated for or dying from cancer or hospice patients or similar conditions as supported by clinical documentation. CA (cancer) or HO (hospice) diag code may be used but store must verify since all scripts will be audited and stores will be liable.	or the preferred drug or a significant potential drug interaction between another drug & the preferred drug(s) exists. Adequate trials include prevention/treatment or common adverse effects associated w/ narcotics (antinausea, antipruritics, etc.) as well as adequate equianalgesic dosing when converting from one narcotic to another. Also, adequate documentation of attempts to titrate dose of preferred agents to achieve adequate pain relief & desired clinical response must be provided. Member's drug regimen for additions &/or discontinuations of medications that may affect absorption &/or metabolism of preferred agents must be monitored. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent w/ controlled substance abuse such as: 1.Frequent or persistent early refills of controlled drugs; 2.Multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel, etc.; 3.Breaches of narcotic contracts with any provider; 4.Failure to comply with patient responsibilities in attached opioid documentation (see PA form) including but not limited to failing to submit to and pass pill counts; 5.Failing to take or pass random drug testing; 6.Failing to provide old records regarding prior use of narcotics; 7.Receiving controlled substances from other prescribers that the provider submitting the PA is unaware of 8.Documented history of substance abuse. Substance abuse evaluations may be required for patients with medical records displaying documented substance abuse or potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycontin. 9.Circumventing MaineCare prior authorization requirements for narcotics by paying cash for affected narcotics (prescribers failed to submit prior authorization prior to cash narcotic scripts being filled by member). 10.Requests for any Brand name controlled substance, considered by authorities to be highly abused and diverted (Oxycontin, Percocet, Tylox, Vicodin, Dilaudid, Ultracet...) with an available AB rated generic equivalent will be denied unless it will be provided in a setting that virtually eliminates the risk of diversion. 11.Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity. Hysingla ER- Concomitant use should be avoided with mixed agonist/antagonist analgesics, partial agonist analgesics, and MAOIs. Verify prior trials and failures or intolerance of preferred treatments Methadone – Established users must have a trial and failure of at least 2 preferred drugs for least 2 weeks. Otherwise they will be allowed 180 days to transition to a preferred product.
NARCOTICS - SELECTED	MC/DEL MC/DEL	TRAMADOL HCL TABS TRAMADOL/APAP TABS	MC/DEL MC MC/DEL MC MC MC MC MC MC MC MC	7 8 8 8 8 8 8 8 8 8 8 9	RYZOLT BUPRENEX SOLN BUTORPHANOL NALBUPHINE HCL SOLN QDOLO SOLN SEGLENTIS ¹ STADOL NS SOLN TRAMADOL ER ULTRACET TABS ¹ ULTRAM ER	Use PA Form# 20420 Use PA form #10300 for PAs over the opiate limit 1. Only available if component ingredients are unavailable.	Preferred drugs from this and other narcotic classes must be tried for at least 2 weeks each and failed due to lack of efficacy or intolerable side effects before non-preferred drugs from this class will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but desired product. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity. Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent with controlled substance abuse such as: 1.frequent or persistent early refills of controlled drugs; 2.multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel; 3.breaches of narcotic contracts with any provider; 4.failure to comply with patient responsibilities in attached opioid documentaion (see PA form) including but not limited to failing to submit to and pass pill counts; 5.failing to take or pass random drug testing; 6.failing to provide old records regarding prior use of narcotics; 7.receiving controlled substances from other prescribers that the provider submitting the PA is unaware of. in Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycontin. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity.

Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 300 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.

However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.

Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.

An MME conversion chart is available at www.mainearepd.org. Click on "General Pharmacy Info."

Please see the Pain Management Policy tab for the complete criteria

MISCELLANEOUS NARCOTICS

NARCOTICS - MISC.						
MC/DEL	ACETAMINOPHEN/CODEINE	MC/DEL	8	ABSTRAL	1. Fentanyl OT loz (Barr)	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E.
MC/DEL	ASPIRIN/CODEINE TABS	MC/DEL	8	APADAZ	and Capital and codeine suspension products require PA for users over 18 years of age. PA is not required if under 18 years of age.	
MC/DEL	BUTAL/ASA/CAFF/COD CAPS	MC/DEL	8	ASCOMP/CODEINE CAPS		
MC	BUTALBITAL/ASPIRIN/CAFFEI CAPS	MC/DEL	8	BUTALBITAL/APAP/CAFFEINE/ CAPS		
MC	CAPITAL AND CODEINE SUSP ¹	MC/DEL	8	BUTALBITAL COMPOUND- CODEINE CAP		
MC	CAPITAL/CODEINE SUSP ¹	MC	8	DEMEROL		
MC/DEL	CODEINE PHOSPHATE SOLN	MC/DEL	8	DILAUDID		Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 300 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.
MC/DEL	CODEINE SULFATE TABS	MC	8	DILAUDID-HP SOLN	2. Oxycodone/acet 10/650 is 8 times more expensive. Use twice as many of oxycod/acet 5/325 instead.	
MC/DEL	ENDOCET TABS ³	MC	8	FENTANYL CITRATE SOLN	You can mix andmatch preferred strengths of oxycodone and oxycodone/acet to minimize acet. dose similar to certain non-preferred drugs.	
MC/DEL	ENDODAN TABS	MC/DEL	8	FENTORA		However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.
MC/DEL	FENTANYL OT LOZ ¹	MC/DEL	8	FIORICET/CODEINE CAPS		
MC/DEL	FENTANYL OT LOZ1	MC	8	FIORINAL/CODEINE #3 CAPS		
MC/DEL	HYDROCODONE/ACETAMINOPHEN	MC	8	FIORTAL/CODEINE CAPS		
MC/DEL	HYDROMORPHONE HCL ³	MC/DEL	8	HYDROCODONE/IBUPROFEN		Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.
MC	LORTAB ELX	MC/DEL	8	HYDROMORPHONE ER		
MC/DEL	MEPERIDINE SOL	MC/DEL	8	HYDROMORPHONE RECTAL SUPP		An MME conversion chart is available at www.mainearepd.org . Click on "General Pharmacy Info."
MC/DEL	NUCYNTA	MC	8	IBUDONE		
MC/DEL	OXYCODONE TAB	MC/DEL	8	LEVORPHANOL TARTRATE TAB		
MC/DEL	OXYCODONE/ACETAMINOPHEN ^{2,3}	MC/DEL	8	LORCET	3. Only preferred manufacturer's products will be available without prior authorization.	
MC/DEL	ROXICET	MC	8	LORTAB		
MC	ROXIPRIN TABS	MC	8	MAXIDONE TABS		
		MC/DEL	8	MEPERIDINE TABS		Please see the Pain Management Policy for the complete criteria
		MC/DEL	8	NORCO TABS		
		MC/DEL	8	ONSOLIS		
		MC/DEL	8	OXECTA		
		MC/DEL	8	OXYCODONE CAP		
		MC/DEL	8	OXYCODONE/APAP 10/650		
		MC/DEL	8	OXYCODONE/APAP 7.5/500		
		MC/DEL	8	PENTAZOCINE/ACET TABS		
		MC/DEL	8	PENTAZOCINE/NALOXONE TABS		
		MC	8	PERCOCET TABS		
		MC	8	PERCOCET TABS		
		MC	8	PHRENILIN W/CAFFEINE/CODE CAPS		
		MC/DEL	8	ROXICET 5/500 TABS		
		MC	8	ROXICODONE TABS		
		MC/DEL	8	ROXYBOND		
		MC	8	SYNALGOS-DC CAPS		
		MC	8	TALACEN TABS		

				MC	8	TREZIX		
				MC	8	TYLENOL/CODEINE #3 TABS		
				MC	8	TYLOX CAPS		
				MC	8	XOLOX	Use PA Form# 20420	
				MC	8	VICODIN		
				MC	8	VICOPROFEN TABS	Use PA form #10300 for PAs over the opiate limit	
				MC	8	ZYDONE TABS		
				MC	9	ACTIQ LPOP		
				MC	9	CONZIP		
				MC	9	OPANA		
OPIOID DEPENDENCE TREATMENTS	MC		SUBOXONE FILM ²					Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		BUPRENORPHINE/NALOXONE TABS ²	MC		BRIXADI	Use PA Form #20100	Members will continue to be required to follow the criteria listed below:
				MC/DEL		BUPRENORPHINE ^{1,2}		1-Induction period for 30 days
				MC		SUBLOCADE	1. Buprenorphine will only be approved for use during pregnancy.	2-Max dose of 32 mg for induction
				MC		ZUBSOLV	2. See Criteria Section	3-Max dose of 24 mg for maintenance
								4-There is not more than one opioid fill in member's drug profile between current fill of buprenorphine and a prior buprenorphine fill within the past 90 days
								5- Should provide evidence of monthly monitoring including random pill counts, urine drug tests and use of Maine Prescription Monitoring Program reports.
								6- Buprenorphine monotherapy is preferred if member is pregnant and dose not > 24 mg day and pregnancy diagnosis is noted on the prescription.
							Use PA form #20200 for Extended Release Buprenorphine	Brixadi and Sublocade: The prescriber can attest (and medical record should document) that: -member has a documented history of opioid use disorder (OUD), -XRB is being used for the treatment of OUD (rather than pain or any other non-FDA approved indication) and -member's total daily dose of sublingual buprenorphine is less than or equal to 24 mg daily. AND at least one of the following is true: -The member's previous use of sublingual buprenorphine has included misuse, overuse, or diversion. -The member is at high risk of overdose (e.g., individuals leaving incarceration or abstinence-based treatment programs; individuals who are unhoused; or those facing potential gaps in care due to delays in care or geographically limited treatment access). -The member has experienced significant medical complications of OUD and/or of injection drug use. Occurrence should be in the last 5 years, or it should be clearly documented that the risk indicated by this infection or complication is ongoing (Examples of medical complications of OUD include: threatened the function of organs or life or limb threatening and required medical and/or surgical therapy. Examples of medical complications of injection drug use include osteomyelitis, endocarditis, renal failure, joint infection or other serious medical complications directly related to OUD.) -The member has treatment-resistant OUD, including those with ongoing illicit substance use in the context of sublingual buprenorphine treatment as documented by positive urine drug screens or other clear objective evidence, and/or further functional decline with explicit documentation of the functional decline. -The member has a significant intolerance of, or documented allergy to, sublingual buprenorphine (either buprenorphine monotherapy or buprenorphine/naloxone combination therapy) that has resulted in the patient's inability to comply with continued treatment using the sublingual product. (A true allergy is usually accompanied by rash, respiratory symptoms, or anaphylaxis. Other complaints such as bad taste, mouth tingling, etc. do not constitute evidence of allergy or significant intolerance. Formulation preference or convenience are not, in and of themselves, indications for using XRB.) -The member is in ongoing treatment with XRB and would like to continue the medication.
OPIOID WITHDRAWAL AGENTS				MC		LUCEMYRA ¹	1. Clinical PA for appropriate approved use and patient has documented contraindication to clonidine Use PA Form#20420	
NARCOTIC ANTAGONISTS								
NARCOTIC - ANTAGONISTS	MC/DEL		NALTREXONE HCL TABS	MC		EVZIO	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC		NALOXONE INJ	MC		OPVEE ²		
	MC		NARCAN NS	MC		KLOXXADO	1. Will only be approved for	

	MC MC MC	NALOXONE SPRAY OTC VIVITROL INJ ZIMHI	MC/DEL	REVIA TABS ¹	side effects experienced with generic that are not described in the literature as occurring with the brand version. 2. For the treatment of adult and pediatric patients 12 years of age and older.
COX 2 / NSAIDS					
COX 2 INHIBITORS - SELECTIVE / HIGHLY SELECTIVE	MC/DEL MC/DEL MC/DEL MC/DEL	CELECOXIB ^{4,5} KETOROLAC TROMETHAMINE ^{2,3,5} NABUMETONE TABS ⁵ MELOXICAM TABS ^{1,5}	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CELEBREX CAPS ^{4,5} MELOXICAM CAPS ⁵ MOBIC ⁵ MOBIC SUSP ⁵ RELAFEN TABS ⁵ QMIIZ ODT VIVLODEX	Use PA Form# 20420 1. Meloxicam has dosing limits allowing one tablet daily of all strengths without PA. 2. Ketorolac Tromethamine is indicated for the short term (up to 5 days) management of moderately severe acute pain that requires analgesic at the opioid level in adults. Not indicated for minor or chronic pain conditions. 3. Ketorolac has dosing limits allowing 24 tablets for a 5 day supply every 30 days. 4. Dosing limits will be set at a maximum of 400mg daily 5. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.
NSAIDS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CHILDRENS IBUPROFEN DICLOFENAC POTASSIUM TABS DICLOFENAC SODIUM TABS DICLOFENAC SODIUM 1% GEL ¹ ETODOLAC FENOPROFEN CALCIUM TABS FLURBIPROFEN TABS IBUPROFEN INDOMETHACIN KETOPROFEN MECLOFENAMATE SODIUM CAPS NAPROSYN SUSP NAPROXEN SUSP NAPROXEN TABS NAPROXEN SODIUM TABS NAPROXEN SODIUM CAPS NAPROXEN DR TBEC	MC MC MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC	ADVIL TABS ANAPROX TABS ANAPROX DS TABS CAMBIA CATAFLAM TABS CHILDRENS ADVIL SUSP CHILD'S IBUPROFEN SUSP CHILDREN'S MOTRIN SUSP CLINORIL TABS DAYPRO TABS DICLIFENAC GEL EC-NAPROSYN TBEC ETODOLAC ER 600MG FELDENE CAPS FLECTOR PATCH IBU-200 INDOCIN	The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use. 1. Dosing limits apply, please see Dosage Consolidation List. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will be granted for other requests based on failure of at least one generic NSAID from at least 3 different NSAID classes as described in the COX-II PA form. DDI: Diclofenac will now be non-preferred and require prior authorization if it is currently being used in combination with Ilescol. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.

	MC/DEL MC/DEL MC/DEL MC/DEL		OXAPROZIN TABS SULINDAC TABS TOLMETIN SODIUM VOLTAREN GEL	MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC MC MC MC MC MC MC	LICART LODINE LOFENA MOTRIN NALFON CAPS NAPRELAN TBCR NAPROSYN TABS NAPROXEN SODIUM TBCR PENNSAID PIROXICAM CAPS PONSTEL CAPS RELAFEN DS SB IBUPROFEN TABS SPRIX TIVORBEX TOLECTIN V-R IBUPROFEN TABS ZORVOLEX		
NSAID - PPI				MC MC/DEL	PREVACID NAPRA-PAC VIMOVO ¹	1. Use a preferred NSAID and PPI separately. Use PA Form# 20420	
RHEUMATOID ARTHRITIS							
RHEUMATOID ARTHRITIS	MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL		ACTEMRA VIALS ACTEMRA SYRINGES AVSOLA AZATHIOPRINE ENBREL ² ENBREL SURECLICK ² KINERET SOLN LEFLUNOMIDE METHOTREXATE ORENCIA SULFASALAZINE TABS SIMPONI PEN SIMPONI AUTOINJECTOR HUMIRA ^{1,2} XELJANZ ^{3,6} XELJANZ XR XELJANZ XR SOL	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC MC MC MC MC MC MC MC	AMJEVITA ARAVA CIMZIA CYLTEZO ENTYVIO HADLIMA HULIO HYDROXYCHLOROQUINE ² HYRIMOZ IDACIO ILARIS ^{1,3,4} INFLECTRA INFLIXIMAB VIAL JYLAMVO KEVZARA OLUMIANT OMVOH OTREXUP RASUVO ⁷ REDITREX REMICADE RENFLIXIS RINVOQ SIMLANDI VELSIPITY YUFLYMA YUSIMRY XATMEP ⁵ ZYMFENTRA	Use PA Form# 20900 1. Dosing limits apply. Please see dose consolidation list. 2. Established users will be grandfathered. 3. Clinical PA is required to establish diagnosis and medical necessity. 4. Verification of age for appropriate indication. 5. Treatment failure or intolerance to other forms of preferred methotrexate 6. See criteria section	See criteria as listed on Rheumatoid Arthritis PA form. Preferred injectable products allowed without PA if trial of a preferred oral agents (azathioprine, hydroxychloroquine, leflunomide, methotrexate, sulfasalazine tabs) are seen in the members drug profile. Dosing limits apply. Xeljanz is limited to adults with moderate to severe RA and UC who have had an inadequate response or intolerance to methotrexate. Should not be used concomitantly with biologic DMARDs or potent Immunosuppressants. Jylamvo will require using preferred methotrexate if unable please provide clinical rational as why inappropriate. Zymfentra: In adults for maintenance treatment of: Moderately to severely active ulcerative colitis following treatment with an infliximab product administered intravenously. Moderately to severely active Crohn's disease following treatment with an infliximab product administered intravenously. DDI: The concomitant use of Xeljanz® XR with biologic DMARDs or potent immunosuppressants such as azathioprine and cyclosporine are not recommended. The concomitant use of Xeljanz® XR with potent CYP3A4 inducers (e.g. rifampin) is not recommended
ALOPECIA AREATA AGENTS							
ALOPECIA AREATA AGENTS				MC MC/DEL	7 8	OLUMIANT LITFULO	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

[Use PA Form# 20420](#)

MISCELLANEOUS ARTHRITIS

ARTHRITIS - MISC.	MC MC		RIDAURA CAPS MYOCHRYSINE SOLN	MC/DEL		ARTHROTEC ¹	1. The individual components of Arthrotec are available without PA. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The individual components of Arthrotec are available without PA.
-------------------	----------	--	----------------------------------	--------	--	------------------------	---	---

LUPUS-SLE

LUPUS-SLE				MC MC MC		BENLYSTA ¹ LUPKYNIS SAPHNELO	Use PA Form# 20420 1. Approvals will require previous trial of corticosteroids, antimalarials, NSAIDS and immunosuppressives.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Lupkynis is a sensitive CYP3A4 substrate. Co-administration with strong or moderate CYP3A4 inhibitors increases voclosporin exposure, which may increase the risk of Lupkynis® adverse reactions. Co-administration of Lupkynis® with strong CYP3A4 inhibitors (e.g. ketoconazole, itraconazole, clarithromycin) is contraindicated. Reduce Lupkynis® dosage when co-administered with moderate CYP3A4 inhibitors (e.g. verapamil, fluconazole, diltiazem)
-----------	--	--	--	----------------	--	---	--	---

PIK3CA-Related Overgrowth Spectrum (PROS)

PIK3CA-Related Overgrowth Spectrum (PROS)				MC		VIJOICE ¹	Use PA Form# 20420 1. PA required to confirm FDA approved indication.	Preferred drugs must be tried and failed, in step-order, due to lack of efficacy (failure to reach target IOP reduction) or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
---	--	--	--	----	--	----------------------	--	--

MIGRAINE THERAPIES

MIGRAINE - ERGOTAMINE DERIVATIVES				MC/DEL MC		D.H.E. 45 SOLN TRUDHESA	Use PA Form# 10110	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
-----------------------------------	--	--	--	--------------	--	----------------------------	------------------------------------	--

MIGRAINE - CARBOXYLIC ACID DERIVATIVES	MC		DIVALPROEX ER TB24	MC		DEPAKOTE ER TB24	Use PA Form# 10110	
--	----	--	--------------------	----	--	------------------	------------------------------------	--

MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)--Tabs/Nasal	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	1 1 1 1 1 1 2	MIGRANAL NASAL SPRAY RELPAX ¹ RIZATRIPTAN ODT RIZATRIPTAN TABS SUMATRIPTAN TABS ¹ ZOLMITRIPTAN TAB ¹ NARATRIPTAN HCI TABS ¹	MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		AMERGE TABS ^{1,2} AXERT TABS ^{1,2} FROVA TABS ^{1,2} IMITREX NASAL SPRAY ¹ IMITREX TABS ^{1,2} MAXALT ^{1,2,3} MAXALT MLT ^{1,2,3} ONZETRA XSAIL ² SUMATRIPTAN NASAL SPRAY ¹ ZOLMITRIPTAN ODT ZOLMITRIPTAN SPRAY ZOMIG TABS ^{1,2} ZOMIG NASAL SPARY ^{1,2} ZOMIG ZMT TBDP ^{1,2}	1. All drugs in this category have dosing limits. Please refer to dose consolidation table. 2. Must fail all preferred products before non-preferred. 3. Established users will be grandfathered Use PA Form# 10110	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Quantity limit exceptions will require ongoing therapy with therapeutic doses of highly effective prophylactic medication as listed on the Triptan PA form.
---	--	---------------------------------	---	--	--	---	--	--

MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)--Injectables	MC MC/DEL MC/DEL		IMITREX CARTRIDGE ¹ SUMATRIPTAN SYRINGE ¹ SUMATRIPTAN PEN INJCTR ¹	MC/DEL MC MC		TOSYMRA ZEMBRACE ¹ IMITREX PEN INJCTR ¹	Use PA Form# 10110 1. Dosing limits apply. Please refer to the dose consolidation table.	
--	------------------------	--	---	--------------------	--	---	---	--

MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)--Combinations				MC/DEL		TREXIMET ^{1,2}	Use PA Form# 10110 1. Dosing limits apply. Please see dose consolidation list.	
---	--	--	--	--------	--	-------------------------	---	--

						2. Use preferred Sumatriptan and Naproxen separately. Treximet only available if component ingredients of sumatriptan and naproxen are unavailable.		
MIGRAINE - MISC.	MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		AIMOVIG ¹ AJOVY ¹ AJOVY AUTO INJCT ¹ EMGALITY SYRINGE ¹ 200mg/ml EMGALITY PEN ¹ NURTEC ODT ² SPASTRIN TABS	MC MC MC/DEL MC/DEL MC MC MC MC/DEL		BELCOMP-PB SUPP ELYXYB MIGRAZONE CAPS MIGERGOT SUP QULIPTA REYVOW ² UBRELVY ² VYEPTI ² ZAVZPRET ²	Use PA Form# 10110 1. See criteria section 2. Dosing limits apply, please see the dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Aimovig, Ajovy and Emgality: The patient is 18 years of age or older AND patient has a diagnosis of episodic migraine (4-14 headache days per month with migraine lasting 4 hours or more) or chronic migraine (≥ 15 headache days per month, of which ≥ 8 are migraine days, for at least 3 months) AND patient has failed or has a contraindication to an adequate trial (≥ 60 days) of at least 2 medications for migraine prophylaxis from at least 2 different classes. Ubrely is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. This is not indicated for the preventive treatment of migraine. Reyvow is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. Reyvow® is not indicated for the preventive treatment of migraine. Zavzpret: The patient must have a documented side effect, allergy, or treatment failure to preferred oral CGRP Inhibitor and two non-preferred oral CGRP Inhibitors. Nurtec ODT will be preferred after 2 adequate trials of at least two preferred triptans
GOUT								
GOUT	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ALLOPURINOL TABS COLCHICINE TAB FEBUXOSTAT TAB PROBENECID TABS PROBENECID/COLCHICINE TABS	MC/DEL MC MC MC/DEL MC MC		COLCHICINE CAP COLCRYS GLOPERBA ULORIC ¹ MITIGARE ZYLOPRIM TABS	Use PA Form# 20420 1. Failure of therapeutic (300mg) dose of Allopurinol (failure define as not being able to get uric acid levels below 6mg/dl) or severe renal disease.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: The concomitant use of Gloperba® and CYP3A4 inhibitors (e.g. clarithromycin, ketoconazole, grapefruit juice, erythromycin, verapamil, etc.) should be avoided due to the potential for serious and life-threatening toxicity.
MISC.								
ACID SPHINGOMYELINASE DEFICIENCY (ASMD)				MC		XENPOZYME ^{1,2}	1.For treatment of non-central nervous system manifestations of acid sphingomyelinase deficiency (ASMD) in adult and pediatric patients 2. Clinical PA required for appropriate diagnosis and clinical parameters.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANESTHETICS - MISC.	MC MC MC		BUPIVACAINE HCL SOLN LIDOCAINE HCL SOLN MARCAINE SOLN	MC MC/DEL MC		SENSORCAINE-MPF SOLN SYNVISC INJ XYLOCAINE SOLN	Use PA Form# 30130	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
COLD AGGLUTININ DISEASE (CAD)				MC		ENJAYMO ¹	1. Indicated to decrease the need for red blood cell transfusion due to hemolysis in adults with cold agglutinin disease (CAD).	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PRIMARY HYPEROXALURIA TYPE 1 (PH1)						OXLUMO ¹ RIVFLOZA	1. PA is required to establish diagnosis and medical	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the

							Use PA Form# 20420	preferred drug(s) exists.
								Rivfloza: The patient has a diagnosis of Primary Hyperoxaluria Type I (PH1) confirmed via genetic testing (identification of alanine: glyoxylate aminotransferase gene (AGXT) mutation) AND urinary oxalate excretion > 0.5mmol/1.73 m2 or urinary oxalate: creatinine ratio is above the upper limit of normal for age AND is at least 9 years of age AND medication is being prescribed by, or in consultation, with a nephrologist or urologist
SICKLE CELL DISEASE	MC/DEL MC		HYDROXYUREA DROXIA	MC MC MC MC MC/DEL		ADAKVEO CASGEVY ^{2,3} ENDAR ¹ LYFGENIA ^{2,3} OXBRYTA ² SIKLOS	1.Evidence of other preferred L-glutamine products utilization and reason for failure. 2. For the treatment of patients ≥ 12 years of age. 3. PA required to confirm FDA approved indication. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: The concomitant use of Oxbryta and strong CYP3A4 inhibitors or fluconazole may increase voxelotor plasma levels and may lead to increased toxicity.
HUTCHINSON- GILFORD PROGERIA SYNDROME (HGPS)				MC		ZOKINVY ^{1,2}	1.In patients 12 months of age and older with a body surface area (BSA) of 0.39m2 and above 2. PA required to confirm FDA approved indication. Use PA Form# 20420	ZOKINVY: To reduce the risk of mortality in Hutchinson-Gilford Progeria Syndrome (HGPS). For the treatment of processing-deficient Progeroid Laminopathies with either: Heterozygous LMNA mutation with progerin-like protein accumulation OR Homozygous or compound heterozygous ZMPSTE24 mutations
VACCINES	MC/DEL MC MC/DEL MC/DEL		ABRYSVO AREXVY GARDASIL 9 SHINGRIX				Use PA Form# 20420	Gardasil 9 will be preferred by MaineCare for ages 19-45 for FDA approved indications. Under the Maine Immunization Program Gardasil 9 is covered under the Vaccine for Children Program for ages 9-18. Please contact 1-800-867-4775 or 207-287-3746 for assistance. Abrysvo will be a preferred vaccine indicated for active immunization for the prevention of lower respiratory tract disease (LRTD) caused by respiratory syncytial virus (RSV) in individuals 60 years of age and older. Active immunization of pregnant individuals at 32 through 36 weeks gestational age for the prevention of LRTD and severe LRTD caused by RSV in infants from birth through 6 months of age. Arexvy will be preferred for active immunization for the prevention of LRTD caused by respiratory syncytial virus (RSV) in individuals 60 years of age and older. SHINGRIX (>= 50yo) is preferred as of 11-20-20 with respective age edit.
APDS				MC		JOENJA ^{1,2,3}	Use PA Form# 20420 1.Clinical PA required for appropriate diagnosis 2. For the treatment of patients 2 years of age and older. 3. Avoid CYP3A drug drug interaction.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ALPHA- MANNOSIDOSIS				MC		LAMZEDE	Use PA Form# 20420 1.Clinical PA required for appropriate diagnosis	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTI-CONVULSANTS								
ANTICONVULSANTS	MC/DEL MC MC/DEL MC/DEL		CARBAMAZEPINE CARBAMAZEPINE ER CAP CARBATROL CP12 CELONTIN CAPS	MC MC MC/DEL MC	8 8 8 8	APTIOM BANZEL BRIVIACT ⁷ CARBAMAZEPINE SUS	Use PA Form# 20420 All non-preferred meds must be used in specified order	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

MC/DEL	CLOBAZAM	MC	8	DEPAKOTE		
MC/DEL	CLONAZEPAM TABS	MC	8	DEPAKOTE ER		
MC	DEPAKOTE SPRINKLES CPSP	MC	8	DIACOMIT	1. Quantity limit. 5/month	
MC/DEL	DIASTAT ¹	MC/DEL	8	DIVALPROEX SODIUM SPRINKLE CAPS	2. Dosing limits apply, please see dose consolidation list.	Approvals will be for patients with a variety of drug-specific FDA-approved indications and for specific conditions supported by at least two published peer-reviewed double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality after recommendation by the DUR Committee and as long as all first line therapies have been tried and failed at full therapeutic doses for adequate durations (at least two weeks).
MC/DEL	DIAZEPAM GEL ¹	MC	8	ELEPSIA XR ¹⁰		
MC/DEL	DILANTIN	MC	8	EPRONTIA SOLN ¹¹		
MC/DEL	DIVALPROEX SODIUM	MC/DEL	8	FELBATOL	3. Dosing limits apply per strength as well as a maximum daily dose of 600mg. Please see dose consolidation list.	
MC	DIVALPROEX SPRINKLE CAP	MC/DEL	8	FELBATOL SUS		
MC/DEL	EPIDIOLEX ⁸	MC/DEL	8	FELBAMATE SUS		
MC/DEL	EPITOL TABS	MC	8	FINTEPLA ⁹		
MC/DEL	ETHOSUXIMIDE SYRP	MC	8	FYCOMPA ²		
MC/DEL	EQUETRO	MC/DEL	8	HORIZANT	4. Adjunctive therapy 17 and older.	*** SEE CHART AT END OF DOCUMENT
MC/DEL	GABAPENTIN ² CAP	MC	8	GRALISE		
MC/DEL	GABAPENTIN ² TAB	MC/DEL	8	KEPPRA TABS	5. Max dose 2400mg	
MC/DEL	GABAPENTIN SOL	MC/DEL	8	KEPPRA SOLN	6. Clinical PA required for appropriate diagnosis	Topamax and Neurontin - Second line therapy for migraine prophalaxis after trial of at least three preferred preventive medications from Group 1 listed on page 2 of the Acute Migraine PA form.
MC/DEL	GABITRIL TABS	MC/DEL	8	KLONOPIN TABS		
MC/DEL	LACOSAMIDE SOL	MC	8	LAMICTAL IR		
MC/DEL	LACOSAMIDE TAB	MC	8	LAMICTAL ODT		
MC	LAMICTAL CHEW	MC/DEL	8	LEVETIRACETAM INJ		All non-preferred meds must be used in specified order.
MC	LAMICTAL XR	MC	8	LIBERVANT	7. Adjunctive therapy in the treatment of partial-onset seizures in patient's ≥16 years of age with epilepsy.	Please use Drug-Drug Interaction PA form #10400 for this combination.
MC/DEL	LAMOTRIGINE ER ODT	MC/DEL	8	LYRICA CR		
MC/DEL	LAMOTRIGINE IR ²	MC/DEL	8	LYRICA SOL ³		
MC/DEL	LEVETIRACETAM SOLN	MC	8	MOTPOLY XR		
MC/DEL	LEVETIRACETAM TABS	MC/DEL	8	MYSOLINE TABS		
MC/DEL	LEVETIRACETAM ER TABS	MC	8	ONFI	8. Epidiolex is for the treatment of seizures associated with Lennox-Gastaut syndrome (LGS), Dravet syndrome (DS) or TS (Tuberous Sclerosis Complex) in patients 1 years of age and older.	Epidiolex Criteria for Lennox-Gastaut syndrome (LGS) and Dravet: a trial of two drugs (clobazam, levetiracetam, valproate derivatives, lamotrigine, topiramate, rufinamide, or felbamate).
MC/DEL	LYRICA ³	MC/DEL	8	OXCARBAZEPINE SUS		Diacomit is for the treatment of seizures associated with Dravet syndrome (DS) in patients 6 months of age and older and wrighing 7kg or more There are no clinical data to support the use of Diacomit® as monotherapy in DS.
MC/DEL	NAYZILAM ¹	MC	8	OXTELLAR XR ⁵		DDI: Concomitant use of Diacomit® with other CNS depressants, including alcohol, may increase the risk of sedation and somnolence. Concomitant use of strong inducers (CYP1A2, CYP3A4, or CYP2C19 inducers, such as rifampin, phenytoin, phenobarbital, and carbamazepine) should be avoided, or dosage adjustments should be made.
MC/DEL	OXCARBAZEPINE	MC/DEL	8	PHENYTEK CAPS		
MC/DEL	PREGABALIN CAPS	MC/DEL	8	POTIGA		
MC/DEL	PHENYTOIN	MC/DEL	8	PREGABALIN (ORAL) SOL		
MC/DEL	PRIMIDONE TABS	MC	8	ROWEEPRA TAB	9. For seizures associated with Dravet syndrome in patients 2 years of age and older	DDI: Avoid concomitant use of Nayzilam® with moderate or strong CYP3A inhibitors.
MC/DEL	QUDEXY XR	MC	8	SABRIL		
MC/DEL	TEGRETOL SUS	MC	8	SEZABY		
MC/DEL	TOPIRAMATE	MC	8	SPRITAM		
MC/DEL	TOPIRAMATE SPRINKLE IR CAPS	MC	8	SYMPAZAN		
MC/DEL	TRILEPTAL SUS	MC/DEL	8	TEGRETOL TAB	10. Adjunctive therapy 12 and older.	Xcopri criteria: History of trials with at least 4 AEDs (2 generic, 2 branded or Uncontrolled seizures on three AEDs; or Uncontrolled on 2 AEDs given along with VNS. Uncontrolled defined as 3 or more TC seizures per year (increases risk of SUDEP); > 6 disabling seizures per year . Any patient who has gone to the ED 2 or more times in the prior 12 months (who has also tried and failed at least 3 other drugs). Ongoing use requires 50 percent reduction in seizure frequency after three months.
MC/DEL	VALPROIC ACID TABS	MC/DEL	8	TIAGABINE		
MC/DEL	VALPROIC ACID SOL	MC	8	TOPAMAX		
MC	VALTOCO ²	MC/DEL	8	TOPIRAMATE ER CAPS		Motpoly XR: pediatric patient weight must be > 50kg and requires multiple preferred medication trials including generic lacosamide
MC/DEL	ZONISAMIDE	MC	8	TOPAMAX SPRINKLE ER CAPS ²		
		MC	8	TOPAMAX SPRINKLE IR CAPS ²		
		MC/DEL	8	TOPIRAMATE SPRINKLE ER CAPS ²		
		MC	8	TROKENDI ^{2,6}		
		MC/DEL	8	VIMPAT ⁴	11. Initial monotherapy for the treatment of partial-onset or primary generalized tonic-clonic seizures in patients 2 years of age and older.	Libervant: For the acute treatment of intermittent, stereotypic episodes of frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern in patients with epilepsy 2 to 5 years of age as long as all preferred therapies have been tried and failed at full therapeutic doses.
		MC/DEL	8	VIMPAT SOL ⁴	Adjunctive therapy for the treatment of partial-onset seizures, primary generalized tonic-clonic seizures, and seizures associated with Lennox Gastaut syndrome in patients 2 years of age and older. The preventive treatment of migraine in patients 12 years and older.	
		MC	8	XCOPRI		
		MC/DEL	8	ZARONTIN SYRP		
		MC/DEL	8	ZARONTIN CAP		
		MC/DEL	8	ZARONTIN SOL		
		MC	8	ZONISADE		
		MC	8	ZTALMY		
		MC/DEL	9	KEPPRA XR		
		MC/DEL	9	NEURONTIN		
		MC/DEL	9	TEGRETOL-XR TB12		

						<p>Will require a step though topiramate.</p> <p>SEE ANTICONVULSANT INDICATION CHART AT THE END OF THIS DOCUMENT</p> <p>M= Monotherapy A= Adjunctive 9= No Evidence</p> <p>The step orders show the relative strength of evidence for use in bi-polar and will guide prior authorization determinations.</p> <p>Step 4 drugs-no PA required.</p>	
					<p>BIPOLAR DISORDER: STEP ORDER</p> <p>M ~ A</p> <p>4 ~ 4 LAMICTAL</p> <p>4 ~ 4 LITHIUM</p> <p>4 ~ 4 CARBAMAZEPINE</p> <p>4 ~ 4 VALPROATE</p> <p>4 ~ 4 ATYPICAL ANTIPSYCHOTICS EXC. CLOZAPINE</p> <p>5 ~ 5 TRILEPTAL</p> <p>9 ~ 6 TOPAMAX</p> <p>9 ~ 7 KEPBRA TABS</p> <p>9 ~ 8 GABITRIL TABS</p> <p>9 ~ 9 NEURONTIN</p>		
					<p>PEDIATRIC BIPOLAR1 DISORDER: STEP ORDER</p> <p>M ~ A</p> <p>(6-18 YEARS WITH OR WITHOUT PSYCHOSIS)</p> <p>4 ~ 4 LITHIUM</p> <p>4 ~ 4 CARBAMAZEPINE</p> <p>4 ~ 4 VALPROATE</p> <p>4 ~ 4</p> <p>ATYPICAL ANTIPSYCHOTICS EXC. CLOZAPINE</p> <p>4 ~ 4 LAMICTAL</p> <p>5 ~ 5 TRILEPTA</p>	<p>Two-step 1 preferred drugs must be tried before Trileptal.</p> <p>The step orders show the relative strength of evidence for use in bi-polar and will guide prior authorization determinations.</p> <p>Step 4 drugs-no PA required.</p>	
ANTI-PARKINSON DRUGS							
PARKINSONS - ANTICHOLINERGICS	MC/DEL MC MC/DEL		BENZTROPINE MESYLATE TABS COGENTIN SOLN TRIHEXYPHENIDYL				Use PA Form# 20420
PARKINSONS - ADENOSINE RECEPTOR ANTAGONIST				MC/DEL		NOURIANZ	<p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Avoid use of Nourianz® with strong CYP3A4 inducers (e.g. carbamazepine, rifampin, phenytoin, St. John's wort).</p> <p>Use PA Form# 20420</p>
PARKINSONS - COMT INHIBITORS				MC/DEL MC MC/DEL		COMTAN TABS ONGENTYS TASMAR TABS	<p>Use PA Form# 20420</p> <p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
PARKINSONS - SELECTED DOPAMIN AGONISTS	MC/DEL MC/DEL		PRAMIPEXOLE ROPINIROLE	MC/DEL MC MC/DEL MC/DEL	5 8 8 8	MIRAPEX TABS ¹ REQUIP TABS MIRAPEX ER NEUPRO PATCH	<p>Use PA Form# 20420</p> <p>1. As of 12/08 users of Mirapex will be grandfathered if diagnosis is Parkinsons.</p> <p>Preferred drug must be tried and failed in step-order due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
PARKINSONS- MAOIS				MC		XADAGO	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered

							on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
							Use PA Form# 20420
PARKINSONS - DOPAMINERGICS/CARBI/ LEVO	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL		AMANTADINE HCLCAPS AMANTADINE HCL TABS BROMOCRIPTINE MESYLATE TABS BROMOCRIPTINE MESYLATE CAPS CARBIDOPA/LEVODOPA TABS ³ CARBIDOPA/LEVODOPA ER CARBIDOPA/LEVO/ENTACAPONE TAB LARODOPA TABS SELEGILINE CAPS HCL SELEGILINE TABS HCL	MC/DEL MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC MC		APOKYN AZILECT ² CARBIDOPA/LEVODOPA RAPDIS ELDEPRYL CAPS GOCOVRI INBRIJA KYNMOBI LODOSYN TABS OSMOLEX ER PARLODEL CAPS PARLODEL TABS RYTARY SINEMET TABS SINEMET TBCR ZELAPAR ¹	1. Approvals will require concurrent therapy with Levodopa and failed trials of Selegiline, Comtan, and Stalevo. 2. Approvals will require trials of Carbidopa/Levodopa, Selegiline, Comtan, and Stalevo. 3. Only preferred manufacturer's products will be available without prior authorization.
				MC/DEL MC		STALEVO ¹ CARBIDOPA/LEVODOPA/ENTACA ¹	Use PA Form# 20420 1.Clinical PA is required to establish diagnosis and medical necessity.
MUSCLE RELAXANTS							
MUSCLE RELAXANTS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL		BACLOFEN TABS CHLORZOXAZONE TABS CYCLOBENZAPRINE HCL 5mg & 10mg TABS LIORESAL INTRATHECAL KIT METHOCARBAMOL TABS TIZANIDINE HCL TABS	MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	7 8 8 8 8 8 8 8 8 8 8 8 8 9 9 9 9	ORPHENADRINE CITRATE CARISOPRODOL 350MG TABS AMRIX DANTRIUM CAPS FLEQSUVY LIORESAL TABS LORZONE LYVISPAH METAXALONE NORFLEX TBCR OZOBAX ROBAXIN-750 TABS VECUROMIUM INJ ZANAFLEX TABS CARISOPRODOL 250MG TABS CHLORZOXAZONE 250mg TABS SKELAXIN TAB SOMA TABS	At least 4 preferred drugs (including tizanidine) must be tried for at least 2 weeks and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an..... acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Elderly patients, over 65, will require written notice of the increased sedative risks and impaired driving.Prior Authorization will not be given for:1. frequent or persistent early refills of controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel, etc. Non-preferred drugs will not be approved if members circumventing MaineCare prior authorization requirements by paying (prescribers failed to submit prior authorization prior to cash narcotic scripts being filled by member). Non-preferred products must be used in specified step order. Lorzone is non preferred and requires at least 4 preferred drugs (including tizanidine) and step care therapy (orphenadrine), as well as reasons for why chlorzoxazone is not acceptable.
				MC/DEL MC/DEL MC MC/DEL MC/DEL MC		CARISOPRODOL/ASPIRIN TABS CARISOPRODOL/ASPIRIN/CODE NORGESIC TABS ORPHENADRINE COMPOUND ORPHENADRINE/ASA/CAFF ORPHENGESIC	Use PA Form# 20420 Individual components are available with PA described in the section above.1. frequent or persistent early refills of non-controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement stolen, dropped in toilet or sink, distant travel, etc.
PARATHYROID HORMONE							
PARATHYROID HORMONE				MC		NATPARA ¹	1. Recommended only for those who cannot be well-controlled on calcium Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the

MC	DAILY MULTI VIT/IRON	MC	FERRALET 90	Click here for the OTC List
MC/DEL	DIALYVITE 1MG	MC	IBERET	
MC/DEL	DIALYVITE 800MG	MC	MATERNA TABS	
MC/DEL	FULL SPECTRUM B	MC	MAXARON	
MC	M.V.I.-12 INJ	MC	MULTIRET FOLIC -500 TBCR	
MC	MULTI-VIT/FLUORIDE	MC/DEL	NATAFORT TABS	
MC/DEL	NATALCARE RX TABS	MC/DEL	NATALCARE CFE 60 TABS ¹	
MC/DEL	NEPHRONEX	MC/DEL	NATALCARE GLOSS TABS ¹	
MC/DEL	NIVA-PLUS (ORAL) TABLET	MC	NATALCARE PIC TABS ¹	
MC/DEL	ONE DAILY TABS	MC	NATALCARE PIC FORTE TABS ¹	
MC/DEL	ONE-DAILY MULTIVITAMINS	MC/DEL	NATALCARE PLUS TABS ¹	
MC/DEL	ONE-TABLET-DAILY	MC	NATALCARE THREE TABS ¹	
MC/DEL	POLY-VIT/IRON/FLUORID SOLN	MC/DEL	NATACHEW CHEW	
MC/DEL	POLY-VITAMIN/FLUORIDE SOLN	MC	NATALFIRST TABS	
MC/DEL	POLY-VITAMINS/IRON SOLN	MC	NATATAB RX TABS	
MC	PRENATA (ORAL) TAB CHEW	MC/DEL	NEPHPLEX RX TABS	
MC/DEL	PRENATAL TABS ¹	MC/DEL	NEPHROCAPS CAPS	
MC/DEL	PRENATAL FORMULA 3 TABS ¹	MC/DEL	NEPHRO-VITE TABS	
MC/DEL	PRENATAL PLUS TABS ¹	MC	NESTABS RX TABS	
MC/DEL	PRENATAL PLUS NF TABS ¹	MC/DEL	NIFEREX	
MC	PRENATAL PLUS/27MG IRON ¹	MC/DEL	OCUVITE TABS	
MC	PRENATAL PLUS/IRON TABS ¹	MC	POLY-VI-FLOR SOLN	
MC	PRENATAL VITAMIN PLUS LOW IRON (ORAL) TAB	MC	POLY-VI-SOL SOLN	
MC/DEL	PRENATAL RX/BETA-CAROTENE ¹	MC	POLY-VI-SOL/IRON SOLN	
MC/DEL	PREPLUS (ORAL) TABLET	MC	POLY-VITAMIN DROPS SOLN	
MC/DEL	RENAL CAPS	MC	PRECARE	
MC/DEL	RENAPHRO CAPS	MC	PREFERA OB	
MC	STRESS TAB NF TABS	MC	PREMESIS RX TABS	
MC	THERAPEUTIC-M TABS	MC	PRENATABS CBF TABS ¹	
MC	THERAVITE LIQD	MC	PRENATAL CARE TABS ¹	
MC/DEL	TRINATAL RX 1 (ORAL) TABLET	MC	PRENATAL MR 90 TBCR ¹	
MC/DEL	TRIVEEN-DUO DHA (ORAL) COMBO. PKG	MC/DEL	PRENATAL MTR/SELENIUM TABS ¹	
MC/DEL	TRI-VITAMIN/FLUORIDE SOLN	MC	PRENATAL OPTIMA ADVANCE TABS ¹	
MC	VITA CON FORTE CAPS	MC	PRENATAL PC 40 TABS ¹	
MC	VITAPLEX PLUS TABS	MC/DEL	PRENATAL RX TABS ¹	
		MC	PRENATE ¹	
		MC	PRENATE ELITE ¹	
		MC	PRIMACARE MISC	
		MC	PROTEGRA CAPS	
		MC	STUARTNATAL PLUS 3 TABS ¹	
		MC	TRI-VI-SOL SOLN	
		MC	TRI-VI-SOL/IRON SOLN	
		MC/DEL	ULTRA NATALCARE TABS	
		MC	ULTRA-NATAL TABS ¹	
		MC	VICON FORTE CAPS	
		MC	VINATAL FORTE TABS ¹	
		MC	VINATE ¹	
		MC/DEL	VINATE ADVANCED TABS ¹	

MISCELLANEOUS MINERALS

MINERALS	MC	CALCARB	MC	ANEMAGEN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
	MC	CALCI-MIX CAPSULE CAPS	MC	CALCET TABS	Please refer to OTC list.	
	MC	CALCIQUID SYRP	MC/DEL	CALCIUM 600-D TABS		
	MC	CALCITRATE/VITAMIN D TABS	MC	CALCIUM/VITAMIN D TABS		
	MC/DEL	CALCIUM	MC	CALTRATE 600 PLUS/VIT D TABS	Click here for the OTC List	
	MC/DEL	CALCIUM CARBONATE	MC	CALTRATE PLUS TABS		
	MC/DEL	CALCIUM CITRATE TABS	MC	CHROMAGEN		

MC/DEL	CALCIUM GLUCONATE TABS	MC	CITRACAL PLUS TABS
MC/DEL	CALCIUM LACTATE TABS	MC	CONTRIN CAPS
MC	CALCIUM/MAGNESIUM TABS	MC	FEOGEN FORTE CAPS
MC/DEL	CALCIUM/VITAMIN D TABS	MC	FEROCON CAPS
MC	CALTRATE 600 TABS	MC/DEL	FERREX 150 CAPS
MC/DEL	CHEWABLE CALCIUM CHEW	MC	FERRO-SEQUELS TBCR
MC	CITRACAL TABS	MC	FE-TINIC CAPS
MC	CITRACAL + D TABS	MC	FE-TINIC 150 FORTE CAPS
MC	CITRUS CALCIUM TABS	MC/DEL	FLUOR-A-DAY SOLN
MC	CITRUS CALCIUM 1500 + D TABS	MC	HEMOCYTE TABS
MC	EFFERVESCENT POTASSIUM TBEF	MC/DEL	K-DUR TBCR
MC/DEL	FEOSTAT CHEW	MC	KLOR-CON PACK
MC	FERATAB TABS	MC	K-LYTE
MC/DEL	FER-GEN-SOL SOLN	MC/DEL	K-PHOS TABS NEUTRAL
MC	FER-IRON SOLN	MC	K-TABS TBCR
MC	FERRONATE TABS	MC	K-VESCENT PACK
MC/DEL	FERROUS SULFATE	MC	MICRO-K 10 MEG CPCR
MC/DEL	FLUOR-A-DAY CHEW	MC	NU-IRON 150 CAPS
MC	FLUORIDE CHEW	MC/DEL	OYSTER SHELL CALCIUM/VITA TABS
MC	FLUORIDE SODIUM CHEW	MC/DEL	POLY-IRON 150 CAPS
MC	FLUORITAB CHEW	MC/DEL	POLYSACCHARIDE IRON CAPS
MC	HM CALCIUM TABS	MC/DEL	POTASSIUM BICARB/CHLORIDE
MC	K+ POTASSIUM PACK	MC/DEL	POTASSIUM CHLORIDE 10MEQ CAPS
MC	KAON ELIX	MC/DEL	POTASSIUM CHLORIDE 8MEQ CAPS
MC	KAON-CL-10 TBCR	MC	TUMS 500 CHEW
MC	KCL 0.075%/D5W/NACL 0.2% SOLN	MC	VIACTIV CHEW
MC	K-EFFERVESCENT TBEF		
MC	KLOR-CON		
MC	KLOTRIX TBCR		
MC/DEL	K-PHOS TABS		
MC/DEL	K-VESCENT TBEF		
MC/DEL	LURIDE CHEW		
MC/DEL	MAGNESIUM GLUCONATE TABS		
MC/DEL	MAGNESIUM SULFATE SOLN		
MC	MAGTABS		
MC	MICRO-K 8 MEG		
MC/DEL	OS-CAL TABS		
MC/DEL	OS-CAL 500 + D TABS		
MC/DEL	OYSCO		
MC/DEL	OYST-CAL TABS		
MC/DEL	OYST-CAL D TABS		
MC/DEL	OYST-CAL/VITAMIN D TABS		
MC/DEL	OYSTER CALCIUM TABS		
MC/DEL	OYSTER SHELL		
MC	PHARMA FLUR		
MC/DEL	PHOSPHA 250 NEUTRAL TABS		
MC	POTASSIUM BICARBONATE TBEF		
MC/DEL	POTASSIUM CHLORIDE 8MEQ		
MC	POTASSIUM EFFERVESCENT		
MC/DEL	SELENIUM TABS		
MC	SLOW-MAG TBCR		
MC/DEL	SODIUM FLUORIDE		
MC	V-R CALCIUM		
MC	V-R OYSTER SHELL CALCIUM		
MC	ZINC SULFATE CAPS		

Please refer to OTC list.

Preferred products that used to require diag codes still require diag codes unless indicated otherwise.

PHENYLKETONURIA (PKU) TREATMENT AGENTS

PHENYLKETONURIA (PKU) TREATMENT AGENTS- INJECTABLES				MC		PALYNZIQ ¹	1. For the treatment of patients ≥ 18 years of age. Use PA Form# 20420	Palynziq is not to be used in combination with Kuvan
PHENYLKETONURIA (PKU) TREATMENT AGENTS- ORAL				MC		KUVAN	Use PA Form# 20420	

MISC. ELECTROLYTES/NUTRITIONALS

ELECTROLYTES/ NUTRITIONALS	MC		INTRALIPID EMUL ¹	MC		BOOST ¹	1. This list of nutritionals is incomplete. All nutritionals still require a PA except for the miscellaneous products listed as preferred. SGA form required for nutritionals unless member has a G/I tube. 2. Formerly known as Omacor. Use PA Form# 20420 & SGA Form	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Medical foods are not to be authorized solely for the purpose of enhancing nutrient intake or managing body weight if the participant is able to eat conventional foods adequately. Medical foods may be approved if the member has a medical condition which precludes or restricts the use of conventional foods and necessitates the use of a formula. Concurrent Stimulant therapy is not an acceptable medical reason/condition for use of medical foods for enhancing nutrient intake or managing body weight. For children under the age of 5, MaineCare will not provide milk- or soy-based standard infant formulas. Regular formulas may be sought through your nearest WIC office. MaineCare will continue to cover medical food for all participants in MaineCare when medical necessity is met. Vascepa requires adjunct therapy for specific indication to reduce TG in those with severe hypertriglyceridemia (500mg per deciliter or more). Proper indication per lab values is required before approval
	MC		P.T.E. -5 SOLN ¹	MC		CASEC POWD ¹		
	MC		SEA-OMEGA CAPS ¹	MC		CHOICE DM LIQD ¹		
				MC		DELIVER 2.0 LIQD ¹		
				MC		DOJOLVI		
				MC		ENFAMIL ¹		
				MC		ENSURE ¹		
				MC		GLUCERNA ¹		
				MC		ISOCAL LIQD ¹		
				MC		KINDERCAL TF LIQD ¹		
				MC		KINDERCAL TF/FIBER LIQD ¹		
				MC		L-CARNITINE CAPS ¹		
				MC		LIPISORB LIQD ¹		
				MC		LOVAZA ^{1,2}		
				MC		MODULEN IBD POWD ¹		
				MC		NUTRAMIGEN POWD ¹		
				MC		NUTREN ¹		
				MC		NUTRITIONAL SUPPLEMENT LIQD ¹		
				MC		NUTRIVENT 1.5 LIQD ¹		
				MC		PEPTAMEN ¹		
		MC	PHENYLADE ¹					
		MC	PHENYL-FREE ¹					
		MC	PKU 3 POWD ¹					
		MC	PREGESTIMIL POWD ¹					
		MC	PROBALANCE LIQD ¹					
		MC	PROSOBEE ¹					
		MC	SCANDISHAKE PACK ¹					
		MC	VASCEPA					

ERYTHROPOEITINS	MC		EPOGEN SOLN	MC	8	ARANESP SOLN ¹	Use PA Form# 10520	Non-Preferred drugs must be tried and failed in step-order, due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please see the EPO PA form for other approval and renewal criteria.
	MC		MIRCERA SYRINGE	MC	8	PROCRIT SOLN ¹	1. Clinical PA is required to establish medical necessity and that appropriate lab monitoring is being done.	
	MC		RETACRIT					

GRANULOCYTE CSF

GRANULOCYTE CSF	MC		NEUPOGEN SYRINGE	MC		FULPHILA	1. Must be used in specified step order.	See approval criteria detailed on Granulocyte Colony Stimulating Factor PA form.
	MC		NEUPOGEN VIAL	MC	8	FYLNETRA		
	MC/DEL		NYVEPRIA SYRINGE	MC	8	GRANIX SYRINGE		
	MC/DEL		ZIEXTENZO	MC	8	GRANIX VIAL		
				MC	8	LEUKINE		
				MC/DEL	8	NIVESTYM		
				MC	8	ROLVEDON		

				MC	8	STIMUFEND		
				MC/DEL	8	ZARXIO		
				MC	9	NEULASTA ¹	Use PA Form# 20520	
GAUCHER DISEASE								
GAUCHER DISEASE				MC		CERDELGA ¹	1. Clinical PA for indication required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA.
				MC		YARGESA ¹		Yargesa: As monotherapy for the treatment of adult patients with mild to moderate type 1 Gaucher disease for whom enzyme replacement therapy is not a therapeutic option (e.g., due to allergy, hypersensitivity, or poor venous access).
							Use PA Form# 20420	
ANTICOAGULANTS / PLATELET AGENTS								
ANTICOAGULANTS	MC		COUMADIN TABS	MC		ARIXTRA SOLN	1. Enoxaparin therapy durations greater than 7 days every 30 days require PA	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA.
	MC/DEL		ENOXAPARIN ¹	MC/DEL		FONDAPARINUX	2. Use other strengths available to obtain desired dose.	
	MC		ELIQUIS	MC/DEL		FRAGMIN INJ	3. Diagnosis required	DDI: Warfarin will require prior authorization if being used in combination with fluconazole, miconazole, or voriconazole.
	MC		ELIQUIS STARTER PACK	MC/DEL		FRAGMIN VIAL	4. For the treatment of patients aged 3 months to less than 12 years of age.	DDI: Warfarin will require prior authorization if being used in conjunction with Gemfibrozil or Fenofibrate.
	MC		HEPARIN SODIUM/NACL 0.9% SOLN	MC/DEL		LOVENOX SOLN		DDI: Rifampin will require prior authorization if being used in combination with Savaysa
	MC		HEP-LOCK SOLN	MC/DEL		LOVENOX 300 ²		
	MC		INNOHEP	MC/DEL		LOVENOX SUBQ SYRINGE		
	MC		HEPARIN LOCK SOLN	MC/DEL		PRADAXA ORAL PELLETS ⁴		
	MC/DEL		HEPARIN LOCK FLUSH SOLN	MC		IPRIVASK		
	MC/DEL		HEPARIN SODIUM SOLN	MC/DEL		SAVAYSAS ³		
	MC/DEL		HEPARIN SODIUM LOCK FLUSH SOLN					
	MC/DEL		PRADAXA					
	MC/DEL		JANTOVEN					
	MC/DEL		WARFARIN SODIUM TABS					
	MC/DEL		XARELTO					
	MC/DEL		XARELTO STARTER PACK					
							Use PA form# 20420	
ANTIHEMOPHILIC AGENTS	MC/DEL		AFSTYLA	MC/DEL		ADYNOVATE VIAL	1. Only if other products unavailable.	Non-preferred will only be approved if other preferred products are unavailable.
	MC		ALPHANATE	MC		ADVATE ^{1,2,5}		
	MC		ALPHANINE SD	MC		ALTUVIIIIO ⁴		Beqvez: FDA Approved Indication: An adeno-associated virus vector-based gene therapy indicated for the treatment of adults with moderate to severe hemophilia B (congenital factor IX deficiency) who:
	MC/DEL		ALPROLIX VIAL	MC/DEL		BEQVEZ	2. Advate may be available with PA in cases of large volume dosing in patients with poor venous access.	· Currently use factor IX prophylaxis therapy, or
	MC/DEL		BEBULIN VIAL	MC/DEL		ESPEROCT		· Have current or historical life-threatening hemorrhage, or
	MC/DEL		BENEFIX SOLR	MC/DEL		ELOCTATE		· Have repeated, serious spontaneous bleeding episodes, and,
	MC/DEL		HELIXATE FS KIT	MC/DEL		HEMGENIX		· Do not have neutralizing antibodies to adeno-associated virus serotype Rh74var (AAVRh74var) capsid as detected by an FDA- approved test.
	MC		HEMLIBRA	MC/DEL		IDELVION		
	MC		HEMOFIL - M	MC/DEL		KOGENATE FS ⁵	3. Not indicated for use in children <12 years of age due to greater risk for hypersensitivity reactions and is not indicated for use in previously untreated patients.	Hemgenix® is an adeno-associated viral vector-based gene therapy for IV infusion after dilution. For treatment of adults with Hemophilia B (congenital Factor IX deficiency) who: Currently use Factor IX prophylaxis therapy, or have current or historical life-threatening hemorrhage, or Have repeated, serious spontaneous bleeding episodes.
	MC		HUMATE-P SOLR	MC/DEL		REBINYN		
	MC/DEL		IXINITY VIAL	MC		RECOMBINATE VIAL ⁵		Altuviiiio is a von Willebrand Factor (VWF) independent recombinant DNA-derived, Factor VIII concentrate indicated for use in adults and children with hemophilia A (congenital factor VIII deficiency) for: Routine prophylaxis to reduce the frequency of bleeding episodes, On-demand treatment and control of bleeding episodes, Perioperative management of bleeding.
	MC/DEL		JIVI ³	MC		ROCTAVIAN ⁴		
	MC		KOATE-DVI	MC		SEVENFACT		
	MC		KONYNE - 80					
	MC/DEL		KOVALTRY					Roctavian: For the treatment of adults with severe hemophilia A (congenital factor VIII deficiency with factor VIII activity <1 IU/dL) without antibodies to adeno-associated virus serotype Inclusion:
	MC		MONARC - M					Severe factor VIII deficiency (less than 1% native factor VIII).
	MC		MONOCLATE - P					Exclusion Criteria:
	MC		MONONINE					Antibodies to the virus AAV5
	MC/DEL		NOVOEIGHT					Factor VIII inhibitors (or history of)
	MC		NOVOSEVEN SOLR					Known significant fibrosis of cirrhosis of the liver, or unexplained elevated LFTs
	MC		NUWIQ					History of inadequate compliance with prophylaxis, or regular bleeds despite adequate prophylaxis
	MC/DEL		PROFILNINE					Conditions in which high-dose steroids are contraindicated.
	MC		RECOMBINATE SOLR					
	MC		REFACTO					

	MC/DEL MC MC/DEL		RIXUBIS VIAL WILATE INJ XYNTHA						Use PA Form# 20420	-Inability to abstain from alcohol for one year Plan to impregnate a partner within 6 months of infusion -Hypersensitivity to mannitol -Active infections, either acute or uncontrolled chronic -HIV infection (limited information on use in this population)	
PLATELET AGGREGATION INHIBITORS	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		ASPIRIN ASPIRIN-DIPYRIDAMOLE ER CPMP 12HR BRILINTA ¹ DIPYRIDAMOLE TABS CLOPIDOGREL 75MG PRASUGREL HCL TAB	MC/DEL MC MC MC/DEL MC/DEL MC/DEL	7 8 8 8 8 8	TICLOPIDINE HCL TABS DURLAZA EFFIENT PERSANTINE TABS PLAVIX TABS ZONTIVITY			Use PA Form# 20715 for Plavix, Effent & Brilinta Use PA form# 20420 for other requests 1. Dosing limits apply, please see dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. A special PA may be obtained at the pharmacy for members scheduled for "stent" placement or have had placement if in the last 12months. Please indicate on prescription date of stent placement. DDI: Plavix will require prior authorization if being used in combination with omeprazole, esomeprazole, cimetidine, fluconazole, ketoconazole, intelence, fluoxetine, ticlopidine, and fluvoxamine. DDI: exists for using maintenance ASA dose >100mg, as it reduces the effectiveness of Brilinta Brilinta- Concomitant use with strong CYP3A4 inhibitors should be avoided (including ketoconazole, itraconazole, atazanavir, and telithromycin). Doses of simvastatin and lovastatin >40mg should be avoided.	
PLATELET AGGR. INHIBITORS / COMBO'S - MISC.	MC/DEL MC/DEL		CILOSTAZOL PENTOXIFYLLINE ER TBCR	MC/DEL MC/DEL MC/DEL MC MC		AGRYLIN CAPS ANAGRELIDE CAPS PLETAL TABS TRENAL TBCR YOSPRALA			Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
HEMATOLOGICALS											
MONOCLONAL ANTIBODY				MC/DEL MC MC/DEL MC MC/DEL MC MC		EMPAVELI ENSPRYNG FABHALTA GAMIFANT SOLIRIS ULTOMIRIS UPLIZNA VOYDEYA			Use PA Form# 20420	A diagnosis of Paroxysmal nocturnal hemoglobinuria (PNH) using the HAM test or flow cytometry is required. In addition, the patient must show evidence of having received a meningitis vaccine at least 2 weeks prior to the start of therapy. Gamifant is recommended for the treatment of adult and pediatric (newborn and older) patients with primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent, or progressive disease or intolerance with conventional HLH therapy. Fabhalta and Ultomiris are recommended for the treatment of adults with paroxysmal nocturnal hemoglobinuria (PNH).	
IMMUNE GLOBULIN	MC MC/DEL MC MC MC/DEL MC/DEL MC		BIVIGAM ¹ CUTAQUIG ¹ GAMUNEX-C GAMMAGARD S-D ¹ HIZENTRA ¹ PANZYGA ¹ PRIVIGEN ¹	MC MC/DEL MC MC/DEL MC MC/DEL		ASCENIV ² CUVITRU GAMMAPLEX INJ HYQVIA OCTAGAM INJ ¹ XEMBIFY			Use PA Form# 20420 1. Clinical PA required 2. For the treatment of patients between 12 to 17 years of age.	Cutaquig is indicated as replacement therapy for primary humoral immunodeficiency (PI) in adults. Xembify is indicated for treatment of primary humoral immunodeficiency (PI) in patients 2 years of age and older. Asceniv indicated for the treatment of primary humoral immunodeficiency (PI) in adults and adolescents (12 to 17 years of age). PI includes but is not limited to the humoral immune defect in congenital agammaglobulinemia, common variable immunodeficiency (CVID), X-linked agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies (SCID).	
HEREDITARY ANGIOEDEMA	MC MC MC MC/DEL	PROPHYLAXIS				PROPHYHLAXIS				1. Clinical PA is required to establish diagnosis and medical necessity. 2. For the treatment of patients ≥ 12 years of age.	Haegarda is indicated for routine prophylaxis to prevent Hereditary Angioedema (HAE) attacks in adolescent and adult patients
		TREATMENT				TREATMENT					
					MC/DEL		KALBITOR VIAL				
	MC/DEL MC MC/DEL		BERINERT KIT ¹ FIRAZYR ¹ RUCONEST VIAL ¹						Use PA Form# 20420		

HEMATOLOGICAL AGENTS- THROMBOPOIETIN RECEPTOR AGONISTS	MC MC		PROMACTA ¹ NPLATE ¹	MC MC/DEL MC/DEL	ALVAIZ DOPTELET MULPLETA	Use PA Form# 20420 1. Clinical PA required. Must see prior trial with insufficient response to corticosteroids and immunoglobulins.	Doptelet and Mulpelta: For the treatment of thrombocytopenia in adults with chronic liver disease who are scheduled to undergo a procedure.
HEMATOLOGICAL AGENTS-IgAN				MC/DEL MC	FILSPARI ¹ TARPEYO	Use PA Form# 20420 1. PA required to confirm FDA approved indication.	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists
ANEMIA- BETA THALASSEMIA				MC MC	REBLOZYL ZYNTEGLO	Use PA Form# 20420	Reblozyl is indicated for the the treatment of anemia in adult patients with beta thalassemia who require regular red blood cell (RBC) transfusion. It is not indicated for use as a substitute for RBC transfusions in patients who require immediate correction of anemia. Zynteglo is indicated for the treatment of adult and pediatric patients with β-thalassemia who require regular red blood cell (RBC) transfusions.
HEMATOLOGIC DISORDER TREATMENT AGENTS				MC/DEL MC	CABLIVI TAVALISSE	Use PA Form# 20420	Tavalisse is recommended for patients at risk of bleeding when one line of therapy (steroids, IVIG, splenectomy) has failed. Cablivi is recommended for the treatment of adult patients with acquired thrombotic thrombocytopenic purpura (aTTP), in combination with plasma exchange and immunosuppressive therapy.
COMPLEMENT RECEPTOR ANTAGONIST				MC	TAVNEOS	Use PA Form# 20420	
WHIM SYNDROME AGENTS				MC	XOLREMDI	Use PA Form#20420	Xolremdi: In patients 12 years of age and older with WHIM syndrome (warts, hypogammaglobulinemia, infections, and myelokathexis) to increase the number of circulating mature neutrophils and lymphocytes.
HEMOSTATIC							
HEMOSTATIC	MC/DEL MC		AMICAR AMINOCAPROIC ACID	MC MC	FIBRYGA RIASTAP	Use PA Form# 20420	Fibryga and Riastap are indicated for the treatment of acute bleeding episodes in adults and adolescents with congenital fibrinogen deficiency, including afibrinogenemia and hypofibrinogenemia. Fibryga® is not indicated for dysfibrinogenemia.
ACUTE HEPATIC PORPHYRIA (AHP)							
ACUTE HEPATIC PORPHYRIA (AHP)				MC	GIVLAARI	Use PA Form# 20420	Givlaari is indicated for the treatment of adults with acute hepatic porphyria (AHP).
PYRUVATE KINASE DEFICIENCY AGENTS							
PYRUVATE KINASE DEFICIENCY AGENTS				MC	PYRUKYND ¹	Use PA Form# 20420 1.PA required to confirm FDA approved indication.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s)
OP. - ANTIBIOTICS							
	MC MC MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL		AK-SPORE OINT BACITRACIN/NEOMYCIN/POLYM BACITRACIN/POLYMYXIN B OINT CHLOROPTIC SOLN ERYTHROMYCIN OINT NEOSPORIN SOLN POLYSPORIN TRIMETHOPRIM SULFATE/POLY TOBRAMYCIN SULFATE SOLN	MC MC MC MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC	AK-POLY-BAC OINT AK-SULF OINT AK-TOB SOLN AZASITE BACITRACIN OINT BLEPH-10 SOLN GATIFLOXACIN DROPS GENTAMICIN SULFATE GENTAK ILOTYCIN OINT LEVOFLOXACIN DROPS NEOMYCIN/BACI/POLYM OINT NEOMYCIN/POLYMYXIN/GRAMIC NEOSPORIN OINT OCUSULF-10 SOLN OCUTRICIN SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

				MC/DEL MC/DEL MC/DEL MC	POLYTRIM DROPS SULFACETAMIDE SODIUM DROPS SULFACETAMIDE SODIUM OINT TERAK OINT		
OP. - ANTI-PARASITIC				MC	XDEMVY ¹	Use PA Form# 20420 1. For the treatment of Demodex biopharitis.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - RHO KINASE INHIBITORS	MC		RHOPRESSA				on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s)
						Use PA Form# 20420	
OP. - QUINOLONES	MC/DEL MC/DEL MC/DEL MC/DEL		CILOXAN OINT CIPROFLOXACIN SOL 0.3% OFLOXACIN QUIXIN SOLN	MC/DEL MC/DEL MC	BESIVANCE CILOXAN SOLN OCUFLOX SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - QUINOLONES-4TH GENERATION	MC/DEL		MOXIFLOXACIN 0.5% SOLN (Generic Vigamox)	MC	ZYMAXID	Use PA Form# 20420	
OP. - ARTIFICIAL TEARS AND LUBRICANTS	MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC MC MC MC		ARTIFICIAL TEARS OINT ARTIFICIAL TEARS SOLN CELLUVISC SOLN EYE LUBRICANT OINT GENTEAL LIQUITEARS SOLN MAJOR TEARS SOLN PURALUBE OINT PURALUBE TEARS SOLN REFRESH SOLN OP REFRESH PLUS SOLN ¹ REFRESH PM OINT	MC/DEL MC MC MC MC/DEL MC MC MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC MC/DEL MC	ARTIFICIAL TEARS SOLN OP BION TEARS SOLN DRY EYES OINT DURATEARS OINT HYPO TEARS ISOPTO TEARS SOLN LACRI-LUBE LUBRIFRESH P.M. OINT MURINE SOLN MUROCEL SOLN NATURE'S TEARS SOLN REFRESH SOLN REFRESH TEARS SOLN ¹ TEARGEN SOLN TEARISOL SOLN TEARS NATURALE TEARS PURE SOLN TEARS RENEWED OINT THERATEARS SOLN V-R ARTIFICIAL TEARS SOLN	Use PA Form# 20420 1. Dosing limits apply, please see dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - BETA - BLOCKERS	MC/DEL MC/DEL MC/DEL MC/DEL		BETOPTIC-S SUSP CARTEOLOL HCL SOLN LEVOBUNOLOL HCL SOLN METIPRANOLOL SOLN	MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL	BETAGAN SOLN BETAXOLOL HCL SOLN ISTALOL OCUPRESS SOLN OPTIPRANOLOL SOLN TIMOPTIC SOLN TIMOLOL DROP TIMOLOL SOL-GEL TIMOPTIC-XE SOLG	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - ANTI-INFLAMMATORY / STEROIDS OPTH.	MC MC/DEL		AK-SPORE HC OINT ALREX SUSP	MC MC	AK-TROL SUSP BAC/POLY/NEOMY/HC OINT	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL	DEXAMETH SOD PHOS SOLN FLAREX SUSP FLUOROMETHOLONE SUSP FML DROPS SUSP 1% FML FORTE SUSP FML S.O.P. OINT LOTEMAX OINT LOTEMAX SUSP LOTEMAX SM DROPS GEL 0.38% NEO/POLY/DEXAMETH OINT NEO/POLY/DEXAMETH SUSP PRED-G SUSP PRED FORTE SUSP 1% PRED MILD SUSP PREDNISOLONE TOBRADEX OINT TOBRADEX SUSP TOBEX OINT SULFACETAMIDE/PREDNISOLONE ZYLET SUSP	MC MC MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC		BLEPHAMIDE S.O.P. OINT BLEPHAMIDE SUSP BROMDAY EFLONE SUSP FLUOR-OP SUSP ILUVIEN IMPLANT INVELTYS LOTEMAX GEL MAXITROL OPTH OINT 0.1% NEO/POLY/BAC/HC OINT NEOM/POLY/DEX OPTH OINT 0.1% OMNIPRED DROPS SUSP OZURDEX PRED-G S.O.P. OINT PREDNISOLONE SODIUM PHOSPHATE SOL RETISERT IMPLANT SULFACET SOD/PRED SOLN TRIESENCE VIAL TOBRADEX ST TOBRAMYCIN SUSP DEXAMETHASONE VASOCIDIN SOLN VEXOL SUSP XIPERE		preferred drug(s) exists.
OP. - PROSTAGLANDINS	MC/DEL MC MC/DEL MC/DEL	LATANOPROST SOL 0.005% LUMIGAN SOLN ROCKLATAN TRAVATAN-Z	MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	7 8 8 8 8 8 8 8 8 8	ZIOPTAN BIMATOPROST 0.03% DROPS DURYSTA IYUZEH RESCULA ^{1,2,3} TRAVATAN SOLN TRAVOPROST VYZULTA XALATAN SOLN ¹ XELPROS	1. All preferreds must be tried. 2. Dosing limits apply, please see dosing consolidation list. 3. Clinical PA is required to establish diagnosis and medical necessity. Use PA Form# 20420	Preferred drugs must be tried and failed, in step-order, due to lack of efficacy (failure to reach target IOP reduction) or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - CYCLOPLEGICS	MC MC/DEL MC/DEL MC/DEL	AK-PENTOLATE SOLN ATROPINE SULFATE CYCLOPENTOLATE HCL SOLN ISOPTO HYOSCINE SOLN	MC/DEL MC MC/DEL MC		CYCLOGYL SOLN ISOPTO ATROPINE SOLN ISOPTO HOMATROPINE SOLN MUROCOLL-2 SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - MIOTICS - DIRECT ACTING	MC/DEL MC MC MC/DEL MC/DEL	ISOPTO CARBACHOL SOLN ISOPTO CARPINE SOLN PILOCAR SOLN PILOCARPINE HCL SOLN PILOPINE HS GEL				Use PA Form# 20420	
OP. - SELECTIVE ALPHA ADRENERGIC AGONISTS	MC MC MC MC/DEL MC/DEL	ALPHAGAN SOLN ALPHAGAN P 0.1% SOLN ALPHAGAN P 0.15% SOLN BRIMONIDINE DROPS 0.2 % SIMBRINZA	MC/DEL MC/DEL		BRIMONIDINE TARTRATE DROPS 0.15 % IOPIDINE SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - ANTI-ALLERGICS	MC/DEL MC MC/DEL MC/DEL MC	AZELASTINE HCL DROPS BEPREVE CROMOLYN SODIUM DROPS KETOTIFEN FUMARATE DROPS LASTACFT	MC MC/DEL MC/DEL MC MC/DEL	8 8 8 8 8	ALOCRIOL SOLN ALOMIDE SOLN EMADINE SOLN OPTICROM SOLN PATANOL SOLN	Use PA Form# 20420	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC/DEL MC/DEL MC/DEL		OLOPATADINE HCL 0.1% OLOPATADINE HCL 0.2% ZADITOR SOLN	MC MC/DEL	8 9	ZERVIAE EPINASTINE			
OP. ANTI-ALLERGICS- MASTCELL STABILIZER CLASS				MC/DEL		ALAMAST SOLN		Use PA Form# 20420	
OP. - CARBONIC ANHYDRASE INHIBITORS/COMBO	MC/DEL MC MC/DEL MC/DEL		AZOPT SUSP COMBIGAN DORZOLAMIDE DORZOLAMIDE/TIMOLOL	MC/DEL		COSOPT SOLN PF		Use PA Form# 20420	
OP. - NSAID'S	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		DUREZOL KETOROLAC OPTH 0.4% KETOROLAC OPTH 0.5% MAXIDEX SUSP NEVANAC PREDNISOLONE DROPS	MC MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC MC/DEL	8 8 8 8 8 8 8 8 8 8 8 8 9	ACULAR LS ¹ ACULAR SOLN ¹ BROMSITE ¹ DEXAMETHASONE DROPS DICLOFENAC OPTH 0.1% FLURBIPROFEN SODIUM SOLN ILEVRO LOTEMAX DROPS GEL SM PROLENSA OCUFEN SOLN ¹ XIBROM ¹ VOLTAREN SOLN ¹ ACUVAIL ¹ BROMFENAC	1. Must fail all preferred products before non-preferred.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Use PA Form# 20420	
OP. - OF INTEREST	MC/DEL MC MC MC		CYCLOSPORINE OPTH 0.05% LUCENTIS RESTASIS DROPPERETTE XIIDRA	MC MC MC MC/DEL MC MC MC MC MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC MC MC MC		BYOOVIZ BEOVU BOTOX SOLR CEQUA CIMERLI CYCLOSPORINE DROPPERETTE CYSTADROPS ¹ CYSTARAN ¹ EYLEA EYLEA HD ¹ EYSUVIS ² IZERVAY ¹ OXERVATE LUCENTIS LUXTURNA MIEBO RESTASIS MULTIDOSE DROPS SUSVIMO SYFOVRE TYRVAYA VABYSMO VERKAZIA VEVYE	1. PA required to confirm appropriate diagnosis and clinical parameters for use. 2. For the short-term (up to two weeks) treatment of the signs and symptoms of dry eye disease.	Must fail adequate trials of multi agents from artificial tears and lubricant category. Beovu is non-preferred and indicated for the treatment of Neovascular (wet) Age-Related Macular Degeneration (AMD) Luxturna will be considered for the treatment of patients with confirmed biallelic RPE65 mutation-associated retinal dystrophy. Patients must have viable retinal cells as determined by the treating physician(s). Vevye - Must fail adequate trials of multi agents from artificial tears and lubricant category and a preferred cyclosporine alternative. Oxervate is non-preferred and is indicated for the treatment of neurotrophic keratits. Eylea is non-preferred and indicated for the treatment of: Neovascular (wet) Age-Related Macular Degeneration (AMD), Macular Edema following Retinal Vein Occlusion (RVO), Diabetic Macular Edema (DME), Diabetic Retinopathy (DR) Miebo is non-preferred and is indicated for the treatment of the signs and symptoms of dry eye disease (DED). Syfovre is non-preferred and is indicated for the treatment of geographic atrophy (GA) secondary to age-related macular degeneration (AMD). Use PA Form# 20420	
DERMATOLOGICAL									
ISOTRETINION, ACNE	MC MC MC MC		AMNESTEEM ¹ CLARAVIS ¹ MYORISAN ¹ ZENATANE ¹	MC MC		ABSORICA ABSORICA LD	1. Users 24 or under, PA will not be required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Use PA Form# 20420	
TOPICAL - ACNE PREPARATIONS	MC MC/DEL MC/DEL		ERYDERM SOLN ERYTHROMYCIN GEL ERYTHROMYCIN SOLN	MC/DEL MC/DEL MC		ADAPALENE 0.3% GEL AKLIEF ⁶ ALTINAC CREA	1. Users 24 or under, PA will not be required. 2. Dosing limits allowing	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	

TOPICAL - ANTIVIRALS				MC/DEL MC/DEL MC MC		ACYCLOVIR OINT DENA VIR CREA ^{1,3} YCANTH ZOVIRAX OINT ^{1,2}	1. Must fail oral treatment with Acyclovir or Valacyclovir. 2. Approvals limited to 1 tube per 180 days. 3. Dosing limits apply, please see dosing consolidation list. 4. For the topical treatment of molluscum contagiosum in adult and pediatric patients 2 years of age and older.	
TOPICAL - ANTINEOPLASTICS	MC		EFUDEX	MC/DEL MC/DEL MC MC/DEL		CARAC CREA FLUOROURACIL SOLARAZE GEL ZYCLARA		Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - BURN PRODUCTS	MC MC/DEL MC MC MC/DEL		FURACIN CREA SILVER SULFADIAZINE CREA SSD AF CREA SSD CREA THERMAZENE CREA	MC/DEL		SILVADENE CREA		Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - CORTICOSTEROIDS	MC MC/DEL MC MC MC MC MC MC MC MC MC/DEL MC MC MC MC MC MC MC/DEL		<p style="text-align: center;">LOW POTENCY</p> DERMA-SMOOTH- FS BODY HYDROCORTISONE CREA HYDROCORTISONE LOTN HYDROCORTISONE LOTN TEXACORT SOLN <p style="text-align: center;">MEDIUM POTENCY</p> DESOXIMETASONE 0.05% CREA/GEL FLUTICASONE PROPIONATE CREA/OINT HYDROCORTISONE BUTYRATE HYDROCORTISONE OINT HYDROCORTISONE VALERATE MOMETASONE FUROATE OINT TRIAMCINOLONE ACETONIDE .025-.1% <p style="text-align: center;">HIGH POTENCY</p> DESONIDE ¹	MC/DEL MC MC MC/DEL MC MC MC MC MC MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL		<p style="text-align: center;">LOW POTENCY</p> ACLOVATE ANUSOL HC-1 OINT DESONATE GEL FLUOCINOLONE ACETONIDE FLUOCINOLONE HALOG HYDROCORTISONE POWD LIDA MANTLE HC CREA PROCTOCORT CREA VERDESO <p style="text-align: center;">MEDIUM POTENCY</p> BESER LOTION ³ CLODERM CREA CORDRAN CUTIVATE CREA / OINT CUTIVATE LOTN DERMATOP ELOCON OINT KENALOG AERS LOCOID LUXIQ FOAM	1. Dosing limits apply, please see dosing consolidation list. 2. Treatment beyond 4 weeks is not recommended. 3. For the treatment of patients ≥ 12 years of age. 4. For the treatment of patients ≥ 18 years of age.	At least 1 drug from each potency of preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC		TRIAMCINOLONE ACETONIDE .5%	MC MC MC MC/DEL MC	PANDEL CREA TOPICORT TOPICORT LP CREA TOVET FOAM ⁹ WESTCORT		
					HIGH POTENCY		
	MC/DEL MC/DEL MC MC		AUGMENTED BETA DIP BETAMETHASONE VALERATE DIFLORASONE DIACETATE HALOBETASOL	MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	AMCINONIDE CREA BETAMETHASONE DIPROPIONATE DESOXIMETASONE 0.25% CREA/OINT		
					VERY HIGH POTENCY		
					VERY HIGH POTENCY		
					BRYHALI LOTN CLOBETASOL PROPINATE LOTN CLOBETASOL PROPINATE SHAMPOO 0.05% CORMAX DIPROLENE IMPEKLO ⁴ LEXETTE OLUX FOAM PSORCON PSORCON E SERNIVO SPRAY ² TEMOVATE ULTRAVATE		
					MISCELLANEOUS		
	MC		PROCTO-KIT CREA 1%	MC/DEL MC/DEL MC/DEL MC MC/DEL MC			
TOPICAL - STEROID LOCAL ANESTHETICS				MC	EPIFOAM FOAM	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - STEROID COMBINATIONS	MC		DERMA-SMOOTH-FS SCALP	MC	CARMOL-HC CREA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - EMOLLIENTS	MC/DEL MC MC		AMMONIUM LACTATE CREA ¹ AMMONIUM LACTATE LOTN 12% ¹ VITAMIN A & D MEDICATED OINT	MC MC MC MC MC	LAC-HYDRIN CREA ¹ LAC-HYDRIN LOTN 12% MEDERMA GEL MIMYX RENOVA CREA	Use PA Form# 20420 1. Dosing limits still apply. Please see dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ENZYMES / KERATOLYTICS / UREA				MC MC MC	CARMOL 40 CREA SALEX CREA SALEX LOTN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Ziox, Panafil and Papain products have been removed from the PDL due to FDA safety concerns regarding drugs containing Papain.
TOPICAL - GENITAL WARTS	MC/DEL		IMIQUIMOD 5% ²	MC/DEL MC/DEL MC/DEL MC MC MC	5 PODOFILOX SOLN 8 CONDYLOX ¹ 8 ALDARA ¹ 8 PICATO 8 VEREGEN ¹ 8 ZYCLARA ¹	Use PA Form# 20420 1. Non-preferred products must be used in specified order. 2. Dosing limits still apply. Please see dose consolidation list.	
TOPICAL - LOCAL ANESTHETICS	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AF CAPSICUM OLEORESIN CREA CAPSAICIN CREA CAPSAICIN PATCH DIBUCAINE OINT ELA-MAX ¹ LIDOCAINE/PRILOCAINE CREA ¹ LIDOCAINE CREAM LIDOCAINE GEL	MC/DEL MC/DEL MC MC MC MC MC MC/DEL	EMLA PADS EMLA CREA LIDA MANTLE CREA LIDODERM PTCH PONTOCAINE SOLN SYNERA ZOSTRIX ZTLIDO ²	1. Lidocaine/Prilocaine cream and Ela-Max products require PA for users over 18 years of age. 2. Dosing limits still apply. Please see dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC/DEL		LIDOCAINE PTCH 5%			CONSOLIDATION LIST	
TOPICAL - DEPIGMENTING AGENTS				MC MC MC MC/DEL MC/DEL MC MC MC	8 8 8 8 8 8 8 9	ALUSTRA CREA EPIQUIN MICRO GLYQUIN CREA HYDROQUINONE CREA HYDROQUINONE/SUNSCREENS SOLAQUIN FORTE CREA TRI-LUMA CREA ELDOQUIN	Use PA Form# 20420 Use PA Form# 20420 As per Medicaid Policy, cosmetic drugs are not covered. Non-cosmetic clinical applications will be considered by prior authorization on a case by case basis.
TOPICAL - SCABICIDES AND PEDICULICIDES	MC/DEL MC MC/DEL MC/DEL MC		ACTICIN CREA LICE KILLING SHAM LICE TREATMENT CREME RINS LIQD PERMETHRIN LOTN NATROBA ¹	MC MC MC/DEL MC MC MC/DEL		ELIMITE CREA EURAX LINDANE MALATHION OVIDE LOTN SPINOSAD SUSP	Use PA Form# 20420 1. Dosing limits apply, please refer to dosage consolidation list. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - WOUND / DECUBITUS CARE				MC MC MC		FILSUVEZ REGRANEX GEL VYJUVEK	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Regranex will be approved for diabetic patients in good control (hgba1c <8), who are not smoking, with a stage III or IV WOCN AND NPUAP lower extremity diabetic ulcer and with an adequate blood supply (T _{cp} 02 >30, ABI>0.7 or ASP> 70), and where the underlying cause has been corrected. The wound must be free of infection and have been previously treated with preferred standard therapies for at least 2 months. Maximum approval for 20 weeks. Vyjuvek: For the treatment of wounds in patients 6 months of age and older with dystrophic epidermolysis bullosa (DEB) with mutation(s) in the collagen type VII alpha 1 chain (COL7A1) gene. Filsuvez: The patient has a diagnosis of dystrophic or junctional epidermolysis bullosa. The patient is at least 6 months old and does not have current evidence or history of squamous cell carcinoma or active infection in the area requiring Filsuvez application. The patient has used standard wound care treatments, including silicone or foam dressings without wound resolution Accuzyme and Ethezyme products have been removed from the PDL due to FDA concerns regarding drugs containing Papain.
TOPICAL - ASTRINGENTS / PROTECTANTS	MC		XERAC AC SOLN	MC MC MC MC		LOWILA BAR MOISTURIN DRY SKIN CREA PROSHIELD PLUS SKIN PROTE CREA SURGILUBE GEL	Use PA Form# 20420 1. Dosing limits apply, please refer to dosage consolidation list. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ANTISEPTICS / DISINFECTANTS	MC/DEL		POVIDONE-IODINE SOLN	MC MC MC MC		BETADINE OINT FORMALYDE-10 AERS IODOSORB LAZERFORMALYDE SOLUTION SOLN	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS EYE							
OP. - EYE	MC MC MC MC MC MC/DEL		AK-DILATE SOLN EYE WASH SOLN NAPHAZOLINE HCL SOLN PHENYLEPHRINE HCL SOLN PONTOCAINE SOLN SODIUM CHLORIDE	MC MC/DEL MC		LENS PLUS REWETTING DROPS MURO 128 NEO-SYNEPHRINE SOLN	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS EAR							
EAR	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL		A/B OTIC SOLN ACETASOL SOLN ACETASOL HC SOLN ACETIC ACID ACETIC ACID/HYDROCORTISON ALLERGEN SOLN CARBAMIDE PEROXIDE 6.5% OTIC SOLN. CIPRO HC SUSP CORTISPORIN-TC SUSP	MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC		ANTIBIOTIC EAR SOLN ANTIBIOTIC EAR SUSP CIPRODEX CIPROFLOXACIN HCL DEBROX SOLN FLOXIN FLUOCINOLONE ACETONIDE OIL DROPS 0.01% OTIPRIO OTOVEL	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL		CORTOMYCIN COLY-MYCIN-S SUSP DERMOTIC EAR DROPS SOLN EAR DROPS RX SOLN EAR WAX REMOVAL DROPS NEOMYCIN/POLYMYXIN/HC OFLOXACIN 0.3% OTIC					
MOUTH ANTISEPTICS								
MOUTH ANTI-INFECTIVES	MC MC/DEL		NILSTAT SUSP NYSTATIN SUSP	MC MC		MYCELEX TROC ORAVIG	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MOUTH ANTISEPTICS	MC/DEL MC/DEL MC MC		CHLORHEXIDINE GLUCONATE LIDOCAINE VISCOUS SOLN TRIAMCINOLONE IN ORABASE PSTE TRIAMCINOLONE ORADENT PSTE	MC MC MC MC		APHTHASOL PSTE ¹ PERIOGARD SOLN ¹ TRIAMCINOLONE ACETONIDE PSTE ¹	Use PA Form# 20420 1. Must fail all preferred products before non-preferred.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DENTAL PRODUCTS								
DENTAL PRODUCTS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ETHEDENT CREA GEL-KAM CONC GEL-KAM GEL 0.4% PHOS FLUR SOLN SF 5000 PLUS CREA SF GEL STANNOUS FLUORIDE ORAL RI CONC	MC/MC MC/DEL MC/DEL MC		APF GEL GEL DENTAGEL GEL PHOS-FLUR GEL THERA-FLUR-N GEL	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ARTIFICIAL SALIVA/STIMULANTS								
ARTIFICIAL SALIVA/STIMULANTS	MC		SALIVA SUBSTITUTE SOLN	MC MC MC		EVOXAC CAPS RADIACARE SOLR SALAGEN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS ANORECTAL								
ANORECTAL - MISC.	MC MC MC/DEL MC/DEL MC/DEL		CORTENEMA ENEM ELA-MAX 5 CREA HYDROCORTISONE ENEM PROCTOSOL HC CREA PROCTOZONE-HC CREA	MC/DEL MC/DEL MC/DEL MC/DEL MC		ANUSOL-HC CREA CORTIFOAM FOAM PROCTOFOAM HC FOAM PROCTO-KIT CREA 2.5% RECTIV OINT	Use PA Form# 20420	
T-CELL ACTIVATION INHIBITOR								
PSORIASIS BIOLOGICALS	MC MC MC MC MC		ENBREL ^{1,5} ENBREL SURECLICK ¹ HUMIRA ^{1,5} OTEZLA TALTZ ²	MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC MC MC MC MC		AMJEVITA BIMZELX ³ COSENTYX ⁴ CYLTEZO HADLIMA HULIO HYRIMOZ IDACIO ILUMYA ³ SKYRIZI SOTYKTU SPEVIGO SILIQ STELARA TREMIFYA YUFLYMA YUSIMRY	1. Dosing limits apply, please refer to dosage consolidation list. 2. Clinical PA required and will be preferred for the indication of plaque psoriasis, psoriatic arthritis and ankylosing spondylitis. 3. For the treatment of adults with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy. 4. Please see criteria section	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Cosentyx approvals for 300mg dose(s) must use "300DOSE" package (containing 2 x 150mg pens or syringes). It is recommended to assess for TB infection prior to starting treatment with Taltz®.

5. Will not require a PA if at least one systemic drug such as methotrexate, cyclosporine, methoxsalen or acitretin is in members drug profile.

[Use PA Form# 20910.](#)

ALTERNATIVE MEDICINES

ALTERNATIVE MEDICINES	MC MC		DIMETHYL SULFOXIDE SOLN MELATONIN	MC/DEL		CO-ENZYME Q-10	Use PA Form# 20420.	Will only be approved for specific conditions supported by at least two double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality.
-----------------------	----------	--	--------------------------------------	--------	--	----------------	-------------------------------------	---

CHELATING AGENTS

CHELATING AGENTS	MC/DEL		CUPRIMINE CAPS	MC MC MC/DEL MC MC/DEL		CLOVIQUE DEPEN TITRATABS TABS EXJADE ¹ SYPRINE TRIENTINE CAPS	Use PA Form# 20420 1. FDA indication of treatment of chronic iron overload due to blood transfusions in membes 2	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Clovique® should be used when continued treatment with penicillamine is no longer possible because of intolerable or life endangering side effects.
------------------	--------	--	----------------	------------------------------------	--	--	---	---

ANTILEPROTIC

ANTILEPROTIC				MC		THALOMID CAPS ¹	1. All PA requests for 150mg dosing will require use of Thalomid 100mg and 50mg capsules. Use PA Form# 20420.	Approved for indications of leprosy, treatment-resistant multiple myeloma and AIDS.
--------------	--	--	--	----	--	----------------------------	--	---

ANTINEOPLASTIC AGENTS

ANTINEOPLASTIC AGENTS - ANTIADNDROGENS	MC/DEL		BICALUTAMIDE	MC/DEL		CASODEX	Use PA Form# 20420.	
ANTINEOPLASTIC AGENTS- LHRH ANALOGS	MC/DEL MC/DEL MC/DEL MC/DEL		LUPRON DEPOTSYPHNGEKIT ¹ LUPRON DEPOT- PED KIT ¹ (1-month) LUPRON DEPOT-PED SYRINGEKIT (3-month) TRIPTODUR VIAL	MC/DEL MC/DEL MC/DEL MC/DEL MC		LUPRON DEPOT SYRINGEKIT FIRMAGON ² SUPPRELIN LA (IMPLANT) KIT TRELSTAR VANTAS ²	1. Dosing limits apply, please refer to dosage consolidation list. 2. PA required to confirm FDA approved indication. Use PA Form# 20420.	
ANTINEOPLASTIC AGENTS - TYROSINE KINASE INHIBITORS				MC MC/DEL MC		SPRYCEL ¹ TYKERB ² GLEEVEC ¹	Use PA Form# 20420. 1. Verification of diagnosis is required. 2. PA required to confirm FDA approved indication and to monitor for potential drug-drug interactions.	
ANTINEOPLASTICS-MISCELLANEOUS	MC MC/DEL MC/DEL		AMIFOSTINE MERCAPTOPYRINE OXALIPLATIN	MC MC/DEL MC/DEL MC MC/DEL MC/DEL		DOCEFREZ ELOXATIN ETHYOL LEUPROLIDE PURINETHOL ZOLINZA	Use PA Form# 20420.	

ANTINEOPLASTICS- MONOCLONAL ANTIBODIES	MC/DEL	TRAZIMERA	MC/DEL	ENHERTU	
			MC/DEL	HERCEPTIN	
			MC/DEL	HERZUMA	
			MC	KANJINTI	
			MC	OGIVRI	
			MC/DEL	ONTRUZANT	Use PA Form# 20420

CANCER

CANCER	MC	ALIMTA	MC	ABECMA	1. PA required to confirm appropriate diagnosis and testing.	All non-preferred: A clinical PA is required to confirm appropriate clinical indication for the individual drug request. Specific to each drug all age, clinical testing requirements, previous step therapies, adjunctive drug therapy requirements, and response without disease progression will be also be evaluated for clinical appropriateness. The standard for the appropriate indication will include the FDA label as well as current NCCN guidelines
	MC/DEL	ANASTROZOLE TABS	MC	AKEEGA		
	MC	ERBITUX	MC	ALECENSA	2. Avoid CYP3A drug drug interaction.	
	MC	IMATINIB MESYLATE	MC/DEL	ALIQOPA ³		
	MC/DEL	LETROZOLE	MC	ALUNBRIG ¹		
	MC	RUXIENCE	MC	ALYMSYS		
	MC/DEL	VIDAZA	MC/DEL	ARIMIDEX	3. Clinical PA required for appropriate diagnosis	Scemblis is for the treatment of adult patients with: Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP), previously treated with two or more
	MC	ZIRABEV	MC	AUGTYRO	4. Re-approval will require documentation of response without disease progression and tolerance to treatment	
			MC	AYVAKIT		
			MC/DEL	AVASTIN	5. Dosing limits apply, please see dosage consolidation list.	
			MC/DEL	BALVERSA	6. Max daily dose of 300mg.	
			MC	BAVENCIO ^{1,8}		
			MC/DEL	BENDEKA ³	7. Monitor liver enzymes periodically and stop treatment upon Grade 3 or higher elevation of liver enzymes approved indication	
			MC/DEL	BESPONS ³		
			MC	BESREMI ¹	8. For patients ≥ 12 years of age	
			MC	BLENREP		
			MC/DEL	BOSULIF	9. For the treatment of patients up to 25 years of age with B-cell acute lymphoblastic leukemia (ALL) that is refractory or in second or later relapse.	
			MC/DEL	BRAFTOVI ¹		
			MC	BREYANZI		
			MC	BRUKINSA		
			MC	CABOMETYX ³		
			MC	CAMCEVI		
			MC/DEL	CALQUENCE ³		
			MC	COMETRIQ ^{3,4,5}		
			MC	COTELLIC		
			MC/DEL	COPIKTRA		
			MC	DARZALEX ³		
			MC/DEL	DAURISMO		
			MC/DEL	ELREXFIO		
			MC/DEL	EMPLICITI(IV) ⁸		
			MC	EPKINLY		
			MC/DEL	ERLEADA		
			MC/DEL	ERIVEDGE		
			MC	EXKIVITY		
			MC	FARYDAK		
			MC/DEL	FEMARA	Use PA Form# 20420	
			MC	FOLOTYN		
			MC	FOTIVDA		
			MC	FRUZAQLA		
			MC	GAVRETO		
			MC/DEL	GILOTRIF ^{4,5}		
			MC/DEL	IBRANCE		
			MC	ICLUSIG ³		
			MC/DEL	IDHIFA ³		
			MC	IMBRUVICA		
			MC	IMDELLTRA		
			MC/DEL	IMFINZI		
			MC/DEL	IMJUDO		

MC	IMLYGIC
MC/DEL	INLYTA
MC/DEL	INREBIC
MC	INQOVI
MC	IWILFIN
MC	JAKAFI
MC	JAYPIRCA ^{1,2}
MC	JEMPERLI
MC/DEL	KEYTRUDA ¹
MC	KIMMTRAK
MC	KISQALI ¹
MC/DEL	KOSELUGO
MC	KRAZATI ³
MC	KYMRIAH ^{3,9}
MC	KYPROLIS ¹
MC	LARTRUVO ¹
MC	LENVIMA
MC/DEL	LIBTAYO ¹
MC	LONSURF
MC/DEL	LORBRENA
MC	LOQTORZI
MC	LUMAKRAS
MC/DEL	LUMOXITI ¹
MC	LUNSUMIO ¹
MC	LYNPARZA ¹
MC	LYTGOBI
MC	NEXAVAR ¹
MC	NERLYNX ³
MC	NINLARO(PO)
MC/DEL	NUBEQA
MC	MARGENZA
MC/DEL	MEKINIST ^{3,4}
MC/DEL	MEKTOVI ¹
MC	MONJUVI
MC/DEL	MYLOTARG ³
MC/DEL	MVASI
MC	ODOMZO ^{1,2,5}
MC	OGSIVEO
MC	OJEMDA
MC	OJJAARA
MC	OMISIRGE
MC	ONUREG
MC/DEL	OPDIVO ³
MC	OPDUALAG
MC	ORGOVYX
MC	ORSERDU ^{2,3}
MC	PADCEV
MC	PEMAZYRE
MC	PEPAXTO
MC	PHESGO
MC/DEL	PIQRAY
MC	POLIVY
MC	POMALYST
MC	PORTRAZZA ³
MC	QINLOCK
MC	RETEVMO

MC	REZLIDHIA
MC/DEL	ROZLYTREK
MC	RUBRACA
MC	RITUXAN
MC	RYBREVANT
MC	RYDAPT
MC	RYLAZE
MC	RYTELO
MC/DEL	SARCLISA
MC	SCEMBLIX ¹
MC/DEL	STIVARGA
MC/DEL	SUTENT ^{1,2}
MC/DEL	SYLATRON
MC	TABRECTA
MC	TALVEY
MC/DEL	TAFINLAR ^{3,4,5,6}
MC	TAZVERIK
MC/DEL	TALZENNA ¹
MC/DEL	TAGRISO
MC	TECARTUS
MC	TECENTRIQ ¹
MC	TEPMETKO
MC/DEL	TIBSOVO ¹
MC	TIVDAK
MC	TRODELVY
MC	TRUSELTIQ
MC/DEL	TRUXIMA
MC/DEL	TRUQAP
MC	TUKYSA
MC	UKONIQ
MC/DEL	VANFLYTA
MC	VEGZELMA
MC	VENCLEXTA ³
MC	VERZENIO ³
MC/DEL	VITRAKVI
MC/DEL	VIZIMPRO ¹
MC	VONJO
MC/DEL	WELIREG
MC/DEL	XALKORI
MC/DEL	XPOVIO
MC/DEL	XOSPATA
MC/DEL	XTANDI
MC/DEL	YERVOY
MC	YESCARTA ³
MC/DEL	ZALTRAP
MC	ZEJULA ¹
MC/DEL	ZELBORAF
MC	ZEPZELCA
MC	ZYDELIG
MC/DEL	ZYKADIA
MC	ZYNLONTA
MC	ZYNYZ ¹
MC	ZYTIGA

IMMUNOSUPPRESSANTS	MC/DEL	CYCLOSPORINE MODIFIED	MC/DEL	CELLCEPT	1. For the treatment of adult	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered
--------------------	--------	-----------------------	--------	----------	-------------------------------	--

	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		GENGRAF CAPS MYCOPHENOLATE MYFORTIC NEORAL SOL RAPAMUNE SANDIMMUNE TACROLIMUS CAPS	MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL		CYCLOSPORINE CAPS CYCLOSPORINE SOL. MODIFIED ENVARUSUS XR NEORAL CAP PROGRAF CAPS REZUROCK ¹ ZORTRESS	and pediatric patients 12 years and older with chronic graft-versus-host disease (chronic GVHD) after failure of at least 2 prior lines of systemic therapy Use PA Form# 20420	on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Cyclosporine will now be non-preferred and require prior authorization if it is currently being used in combination with either Lipitor (doses greater than 20mg/day), Crestor, or lovastatin (doses greater than 20mg). DDI: Cyclosporine will require prior authorization when used with Livalo. DDI: All preferred immunosuppressants will require clinical PA for patients over 60 that are currently on fluoroquinolone therapy.
IMMUNOSUPPRESSANTS- Misc.				MC		HYFTOR ^{1,2}	1. For the treatment of patients ≥ 6 years of age. 2. Clinical PA required for appropriate diagnosis and clinical parameters. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PURINE ANALOG								
PURINE ANALOG	MC MC/DEL		AZASAN TABS AZATHIOPRINE TABS	MC/DEL		IMURAN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
K REMOVING RESINS								
K REMOVING RESINS	MC/DEL MC/DEL		LOKELMA SODIUM POLYSTYRENE SULFON	MC/DEL MC/DEL MC		SPS SUSP SPS 30GM/120ML ENEMA SUSP VELTASSA	Use PA Form# 20420	

New drugs are initially non-preferred until reviewed by the DUR Committee and the State. According to State policy, any drug requiring specific diagnosis still requires the specific diagnosis unless otherwise noted within this document.

Last update 01/17

PDL DOSAGE CONSOLIDATION LIST

Tabs/Caps/Patches: Quantities in units

Shaded areas are non-preferred agents - Quantities of these

Sprays/Inhalers/Nebulizers: Quantities in GM, ML, OR MCG

non-preferred agents are available up the limit only with

Injectibles: Quantities in ML

prior authorization

Drug Name	Strength	Limit/Day	Limit/Days	Drug Name	Strength	Limit/Day	Limit/Days
ABILIFY SOLUTION	1MG/ML	30ML	1020/34	ATROVENT HFA	17MCG	12 INHALATIONS	25.8/34
ACCUPRIL	5MG	1	35/35	ATROVENT 30ML	0.03%	12 SPRAYS	30/30
ACCUPRIL	10MG	1	35/35	ATROVENT 15ML	0.06%	16 SPRAYS	45/30
ACCUPRIL	20MG	1	35/35	AVANDIA	2MG	1.5	53/35
ACEON	2MG	1	35/35	AVANDIA	4MG	1	35/35
ACEON	4MG	1	35/35	AVAPRO	75MG	1.5	53/35
ACTONEL	5MG	1	35/35	AVAPRO	150MG	1	35/35
ACTONEL	35MG	1/WK	5/35	AXERT (Step 8)	6.25MG		12/30
ACTOS	All Strengths	1	35/35	AXERT (Step 8)	12.5MG		12/30
ADDERALL XR	5MG	3	90/30	AZELEX	20%		1 TUBE/18
ADDERALL XR	10MG	3	90/30	AZILECT	All Strengths	1	35/35
ADDERALL XR	15MG	3	90/30	BACTROBAN CREAM			1 TUBE/30
ADDERALL XR	20MG	2	60/30	BECONASE AQ	42MCG	8 INHALATIONS	50/30
ADDERALL XR	30MG	1	35/35	BENICAR-HCT	All Strengths	1	30/30
ADEMPAS	All Strengths	1	35/35	BENAZEPRIL	5MG	1	35/35
ADVAIR DISKUS	All Strengths	2	60/30	BENAZEPRIL	10MG	1.5	53/35
ADVAIR HFA	All Strengths	4	120/30	BENAZEPRIL	20MG	1	35/35
ADZENYS XR	All Strengths	1	30/30	BENAZEP/HCTZ	5-6.25	1	35/35
AEROBID	250MCG	8 INHALATIONS	21/35	BENAZEP/HCTZ	10/12.5	1	35/35
AEROBID-M	250MCG	8 INHALATIONS	21/35	BEVESPI AERO		4 INHALATIONS	120/30
ALAVERT-NON DROW	TAB	1	96/96	BONIVA	2.5MG	1	35/35
ALENDRONATE	All Strengths	1/WK	35/35	BOTOX (ADULTS)	100U/ML	1 session/90 days	600U/90
ALTABAX	5GM		1 TUBE/30	BOTOX (CHILDREN>12)	100U/ML	1 session/90 days	400U/90
ALTABAX	15GM		1 TUBE/30	BREO ELLIPTA	100/25MCG	1 INHALATIONS	60/60
ALTABAX	30GM		1 TUBE/30	BRILINTA	All Strengths	2	70/35
ALTACE	1.25MG	1	35/35	BRINTELLIX	All Strengths	1	35/35
ALTACE	2.5MG	1	35/35	BUTRANS		1 patch/WK	4/28
ALTACE	5MG	1	35/35	BYETTA	5mcg inj	0.04ML	1.2ML/30
AMARYL	1MG	1	35/35	BYETTA	10mcg inj	0.08ML	2.4ML/30
AMARYL	2MG	1	35/35	CALAN SR	120MG	1	35/35
AMBIEN	5MG		12/34	CALAN SR	180MG	2	70/35
AMBIEN	10MG		12/34	CALAN SR	240MG	2	70/35
AMBIEN CR	6.25MG		12/34	CARDIZEM CD	120MG/24	1	35/35
AMBIEN CR	12.5MG		12/34	CARDIZEM CD	180MG/24	1	35/35
AMERGE (Step 8)	1MG		12/30	CARDIZEM CD	240MG/24	1	35/35
AMERGE (Step 8)	2.5MG	2.5MG	12/30	CARDIZEM CD	300MG/24	1	35/35
AMLODIPINE	2.5MG	1.5	53/35 DAYS	CARDIZEM CD	360MG/24	1	35/35
AMLODIPINE	5MG	1.5	53/35 DAYS	CARDIZEM LA	120MG/24	1	35/35
AMMONIUM LACTATE CREA	12%		1 TUBE/10	CARDIZEM LA	180MG/24	1	35/35
AMMONIUM LACTATE LOTN	12%		1TUBE/8	CARDIZEM LA	240MG/24	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	5MG	3	90/30	CARDIZEM LA	300MG/24	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	10MG	3	90/30	CARDIZEM LA	360MG/24	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	15MG	3	90/30	CARDURA	1MG	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	20MG	2	60/30	CARDURA	2MG	1.5	53/35
AMPHETAMINE/DEXTROAMPHET ER	30MG	1	90/90	CARDURA	4MG	1.5	53/35
AMPHETAMINE SALT	5,10,15MG	3	105/35	CARTIA XT	120MG	1	90/90
AMPHETAMINE SALT	20MG	2	70/35	CARTIA XT	180MG	1	90/90
AMPHETAMINE SALT	30MG	1	35/35	CARTIA XT	240MG	1	90/90
ANDRODERM	2.5MG	2	60/30	CARTIA XT	300MG	1	90/90
ANDRODERM	5MG	1	30/30	CATAPRES-TTS1	0.1 MG/24HR		5/35
ARAVA	10MG	1	35/35	CATAPRES-TTS2	0.2 MG/24HR		5/35
ARCAPTA	75MCG	1 INHALATION	35/35	CATAPRES-TTS3	0.3 MG/24HR		5/35
ARICEPT	5MG	1	35/35	CEFIXIME	400MG	2	2/7
ARICEPT	10MG	1	35/35	CELEBREX	100MG	1	35/35
ARIPIRAZOLE	2MG	2	180/90	CELEBREX	200MG	2	70/35
ARIPIRAZOLE	5MG	2	180/90	CELEBREX	400MG	1	35/35
ARIPIRAZOLE	10MG	2	180/90	CELEXA	20mg	0.5	17/34
ARIPIRAZOLE	15MG	2	180/90	CELEXA	40mg	1	51/34
ARIPIRAZOLE	20MG	1.5	135/90	CITALOPRAM	10MG	2	180/90
ARIPIRAZOLE	30MG	1	90/90	CITALOPRAM	20MG	2	180/90
ARIXTRA INJECTION	2.5MG/0.5ML		7/30	CITALOPRAM	40MG	1	90/90
ARIXTRA INJECTION	5MG/0.4ML		7/30	CLARINEX	REDI TAB	1	35/35
ARIXTRA INJECTION	7.5MG/0.6ML		7/30	CLEOCIN-T		1 PACKAGE	1/30
ARIXTRA INJECTION	10MG/0.8ML		7/30	CLINDAMYCIN PHOSPHATE		1 PACKAGE	1/30
ARMONAIR	All Strengths	1 INHALATION	60U/30	COMBIVENT	103-18MCG	12 INHALATIONS	30/35
ASMANEX 30 UNITS	220MCG	1 INHALATION	30U/30	Drug Name	Strength	Limit/Day	Limit/Days
ASMANEX 60 UNITS	220MCG	2 INHALATIONS	60U/30	EFFEXOR XR	37.5MG	1	35/35
ASMANEX 120 UNITS	220MCG	4 INHALATIONS	120U/30	EFFEXOR XR	75MG	1	35/35
ATACAND	4MG	1.5	53/35	EMSAM	All Strengths	1	34/34
ATACAND	8MG	1.5	53/35	ENALAPRIL	2.5	1	90/90

ATACAND	16MG	1	35/35
ATRIPLA	600MG	1	35/35
Drug Name	Strength	Limit/Day	Limit/Days
COMETRIQ	80MG	1	35/35
COMETRIQ	20MG	3	105/35
CONCERTA	18MG	1	30/30
CONCERTA	27MG	1	30/30
CONCERTA	36MG	2	60/30
COPAXONE INJ	20MG		1/32
COPAXONE KIT	20MG/ML		1/30
COREG CR	All Strengths	1	34/34
COSENTYX	150MG	1	1/30
CRESTOR	5MG	1	35/35
CRESTOR	10MG	1	35/35
CRESTOR	20MG	1	35/35
CRESTOR	40MG	1	35/35
CYMBALTA	All Strengths	1	35/35
DALMANE	15MG		10/30
DALMANE	30MG		10/30
DAYPRO	600MG	2	70/35
DAYTRANA	10mg/9hr (27.5mg)	1	34/34
DAYTRANA	15mg/9hr (41.3mg)	1	34/34
DAYTRANA	20mg/9hr (55.0mg)	1	34/34
DAYTRANA	30mg/9hr (82.5mg)	1	34/34
DDAVP	5ML		15/34
DENAVIR CREAM			2gm/30
DEPO-PROVERA	150MG/ML		1/90
DEPO-PROVERA	400MG/ML		2.5/90
DEPO-TESTOSTERONE	200MG/ML		20/90
DESMOPRESSIN	0.1MG	12	420/35
DESMOPRESSIN	0.2MG	6	210/35
DESONIDE	0.05%		2 TUBES/30
DESOWEN	0.05%		2 TUBES/30
DETROL LA	2MG	1	35/35
DEXEDRINE	All Strengths	3	90/30
DEXILANT	All Strengths	1	35/35
DEXTROAMPHETAMINE	All Strengths	3	90/30
DICLOFENAC 1% GEL	1% GEL		2 TUBES/30
DIFLUCAN	150MG		1/7
DILACOR XR	240MG/24	1	35/35
DILACOR XR	120MG/24	1	35/35
DILACOR XR	180MG/24	1	35/35
DILTIA - XT	120MG/24	1	90/90
DILTIA - XT	180MG	1	90/90
DILTIA - XT	240MG/24	1	90/90
DILTIAZEM CAP ER	120MG	1	90/90
DILTIAZEM CAP XR	120MG	1	90/90
DILTIAZEM CAP	120MG/24	1	90/90
DILTIAZEM CAP	180MG/24	1	90/90
DILTIAZEM CAP ER	240MG	1	90/90
DILTIAZEM CAP XR	240MG	1	90/90
DILTIAZEM XR CAP	240MG/24	1	90/90
DILTIAZEM CAP	240MG/24	1	90/90
DILTIAZEM CAP	300MG/24	1	90/90
DILTIAZEM CAP	360MG/24	1	90/90
DIOVAN	80MG	1	35/35
DIOVAN - HCT	80 - 12.5	1	35/35
DITROPAN XL	5MG	1	35/35
DITROPAN XL	10MG	2	70/35
DORAL	7.5MG		10/30
DOXAZOSIN	1MG	1	90/90
DOXAZOSIN	2MG	1.5	135/90
DOXAZOSIN	4MG	1.5	135/90
DRYSOL SOL	20%		1 BOTTLE/30DAYS
DURAGESIC PATCHES	12.5MCG/HR		11/33
DURAGESIC PATCHES	25MCG/HR		11/33
DURAGESIC PATCHES	50MCG/HR		11/33
DURAGESIC PATCHES	75MCG/HR		11/33
DURAGESIC PATCHES	100MCG/HR		22/33
DULOXETINE	20MG	3	270/90
DULOXETINE	30MG	3	270/90
DULOXETINE	60MG	2	180/90
EDEX	All Strengths		1/30
Drug Name	Strength	Limit/Day	Limit/Days
ILARIS			2/28

ENALAPRIL	5MG	1.5	135/90
ENALAPRIL	10MG	1.5	135/90
ENALAPR/HCTZ	5-12.5	1	90/90
ENBREL	25MG/ML		8/28
ENBREL SURECLICK			8/28
ESTAZOLAM	1MG		10/30
ESTAZOLAM	2MG		10/30
ESTRING MIS	2MG		1/90
EVENITY		12 DOSES/LIFETIME	12 DOSES/LIFETIME
EVOTAZ	All Strengths	1	30/30
FELODIPINE	2.5MG	1	90/90
FELODIPINE	5MG	1.5	135/90
FENTANYL	25MCG/HR		11/33
FENTANYL	50MCG/HR		11/33
FENTANYL	75MCG/HR		11/33
FENTANYL	100MCG/HR		22/33
FETZIMA	All Strengths	1	35/35
FINASTERIDE	5MG	1	90/90
FLONASE	50MCG	4 SPRAYS	32/34
FLOVENT HFA 44MCG	44MCG	4 INHALATIONS	10.6/30
FLOVENT HFA 110MCG	110MCG	4 INHALATIONS	12/30
FLOVENT HFA 220MCG	220MCG	8 INHALATIONS	24/30
FLOVENT DISKUS	50MCG, 100MCG	4 INHALATIONS	60/30
FLOVENT DISKUS	250MCG	3 INHALATIONS	120/30
FLUCONAZOLE	150MG		1/7
FLUNISOLIDE SOLN	0.025%	16 SPRAYS	75/30
FLUOXETINE CAP	40MG	2	180/90
FLUOXETINE CAP	20MG	4	360/90
FLUOXETINE CAP	10MG	3	270/90
FLURAZEPAM	15MG		10/30
FLURAZEPAM	30MG		10/30
FLUTICASONE SPR		4 SPRAYS	48/90
FLUVOXAMINE	25MG	3	270/90
FLUVOXAMINE	50MG	3	270/90
FOCALIN	All Strengths	3	105/35
FOCALIN XR	All Strengths	1	35/35
FORFIVO XL	All Strengths	1	35/35
FOSAMAX	5MG	1	35/35
FOSAMAX	10MG	1	35/35
FOSAMAX	70MG	1/WK	5/35
FOSAMAX	40MG	2/WK	10/35
FOSINOPRIL	10MG	1.5	135/90
FOSINOPRIL	20MG	2	180/90
FRAGMIN INJ	10000U/ML	2ML	14/7
FRAGMIN INJ	2500U/.2ML	0.4ML	2.80/7
FRAGMIN INJ	25000U/ML	0.8ML	5.6/7
FRAGMIN INJ	5000U/.2ML	0.4ML	2.80/7
FRAGMIN INJ	7500U/.3ML	0.6ML	4.2/7
FROVA TAB (Step 8)	2.5MG		12/30
FULYZAQ	125MG	2	70/35
FUZEON	KIT	1	1/30
FYCOMPA	All Strengths	1	35/35
GABAPENTIN	300MG	9	810/90
GABAPENTIN	400MG	9	810/90
GABAPENTIN	600MG	6	540/90
GABAPENTIN	800MG	4	360/90
GEODON	20MG	2	70/35
GEODON	40MG	2	70/35
GEODON	60MG	2	70/35
GEODON	80MG	2	70/35
GEODON	INJ	2	70/35
GILOTRIF	All Strengths	1	35/35
GLIMEPIRIDE	1MG	1	90/90
GLIMEPIRIDE	2MG	1	90/90
GLUCOSE TES STRP		12	420/35
GLUCAGEN INJ. HYPOKIT			2/30
GLYCOLAX*	255GM		255GM/90
* Available for once daily dosing to members under the age of 18 years			
Drug Name	Strength	Limit/Day	Limit/Days
LUNESTA	2MG		12/34
LUNESTA	3MG		12/34
LUPRON DEPOT INJ	11.25MG	KIT	1/90
LUPRON DEPOT INJ	22.5	KIT	1/90
LUPRON DEPOT INJ	30MG		1/90

HALCION	0.125MG		10/35
HALCION	0.25		10/35
HUMIRA	40mg/0.8ml		4/28
HYDROXYZINE TAB	All Strengths	3	270/90
HYTRIN	1MG	1	35/35
HYTRIN	5MG	1	35/35
HYZAAR	50-12.5	1	35/35
IMDUR	30MG	1.5	53/35
IMDUR	60MG	1.5	53/35
IMITREX (step 8)	25MG		12/30
IMITREX (step 8)	50MG		12/30
IMITREX (step 8)	100MG		12/30
IMITREX VIAL	All Strengths		6 boxes/30
IMITREX CARTRIDGE	All Strengths		12/30
IMITREX NASAL SPRAY	All Strengths		12/30
IMITREX PEN INJCTR	All Strengths		12/30
IMIQUIMOD	5%		12/30
IMIQUIMOD	5%		12/30
INTAL	800MCG	8 INHALATIONS	28.4/34
INVOKANA	All Strengths	1	35/35
IPRATROPIUM 30ML	0.03%	12 SPRAYS	90/90
IPRATROPIUM 15ML	0.06%	16 SPRAYS	135/90
ISOPTIN SR	180MG	2	70/35
IRBESARTAN	All Strengths	1	90/90
ISOPTIN SR	240MG	2	70/35
ISOSORBIDE MONO	30MG	2	180/90
ISOSORBIDE MONO	60 MG	1.5	135/90
JANUMET	All Strengths	2	70/35
JANUVIA	All Strengths	1	35/35
JUVISYNC	All Strengths	1	35/35
KETOPROFEN	100MG	2	180/90
KETOPROFEN	200MG	1	90/90
KETOROLAC	10MG	4.8	24/30
KHEDEZLA	All Strengths	1	35/35
LAC-HYDRIN CREAM	12%		1TUBE/30
LAMICTAL	25MG	6	210/35
LAMICTAL	25MG CHW	6	210/35
LAMICTAL	100MG	2	70/35
LAMISIL	250MG	1	35/35
LAMOTRIGINE	25MG	6	540/90
LAMOTRIGINE	100MG	2	180/90
LANSOPRAZOLE CAPS	All Strengths	1	90/90
LATUDA	All Strengths	1	17/34
LESCOL	20MG	1	35/35
LEVAQUIN	250MG	1	35/35
LEXAPRO	5MG	0.5	15/30
LIPITOR	10MG	1	35/35
LIPITOR	20MG	1	35/35
LIPITOR	40MG	1.5	53/35
LISINOP/HCTZ	10/12.5MG	1	90/90
LINEZOLID	600mg		14/60
LOSARTAN	All Strengths	1	90/90
LOSARTAN- HCT	All Strengths	1	90/90
LOTENSIN	5MG	1	35/35
LOTENSIN	10MG	1.5	35/35
LOTENSIN	20MG	1	53/35
LOTENSIN - HCT	5 - 6.25	1	35/35
LOTENSIN - HCT	10 - 12.5	1	35/35
LOVASTATIN	10MG	1.5	135/90
LOVASTATIN	20MG	1.5	135/90
LOVENOX INJ	30MG/.3ML	0.6	14 injections/7
LOVENOX INJ	40MG/.4ML	0.8	14 injections/7
LOVENOX INJ	60MG/.6ML	1.2	14 injections/7
LOVENOX INJ	80MG/.8ML	1.6	14 injections/7
LOVENOX INJ	100MG/ML	2	14 injections/7
LOVENOX INJ	120MG/.8ML	1.6	14 injections/7
LOVENOX INJ	150MG/ML	2	14 injections/7
LUNESTA	1MG		12/34
Drug Name	Strength	Limit/Day	Limit/Days
NIFEDIPINE ER	90MG	1	90/90
NIFEDIPINE ER,CR	30MG	1	90/90
NORVASC	2.5MG	1.5	53/35 DAYS
NORVASC	5MG	1.5	53/35 DAYS
NURTEC ODT	All Strengths		8/30
NUVARING		1/MO	1/28

LUPRON DEPOT INJ	30MG	KIT	1/90
LYRICA	25,50,75MG	3	102/35
LYRICA	100,150,200MG	3	102/35
LYRICA	225,300MG	2	70/35
MAVIK	1MG	1	35/35
MAVIK	2MG	1	35/35
MAXAIR AUTO	200MCG	12 INHALATIONS	14/30
MAXALT (step 8)	5MG		12/30
MAXALT (step 8)	10MG		12/30
MAXALT MLT (step 1)	5MG		12/30
MAXALT MLT (step 1)	10MG		12/30
MEDROXYPR AC	150MG/ML		1/90
MELOXICAM TABS	All Strengths	1	90/90
METADATE ER	10,20MG	3	90/30
METFORMIN ER	500MG	4	360/90
METHYLIN	All Strengths	3	90/30
METHYLPHENIDATE ER	36mg	2	180/90
METHYLPHENIDATE	All Strengths	3	90/30
METROCREAM		1 PACKAGE	1/30
METROGEL		1 PACKAGE	1/30
METROLOTION		1 PACKAGE	1/30
METRONIDAZOLE CREAM		1 PACKAGE	1/30
METRONIDAZOLE GEL		1 PACKAGE	1/30
METRONIDAZOLE LOTION		1 PACKAGE	1/30
MEVACOR	10MG	1.5	53/35
MEVACOR	20MG	1.5	53/35
MIACALCIN		3.75ml	1 bottle/34
MICARDIS	All Strengths	1	30/30
MICARDIS-HCT	All Strengths	1	30/30
MIGRANAL NASAL SPRAY	All Strengths		12/30
MIRALAX	255G	8.5G	1 bottle/30
MIRALAX	17G/PACKET	0.5 packet	15 packets/30
MIRTAZAPINE	15mg	3	270/90
MOBIC	7.5 MG	1	35/35
MOBIC	15MG	1	35/35
MOEXIPRIL	7.5	1.5	135/90
MONOPRIL	10MG	1.5	53/35
MONOPRIL	20MG	2	70/35
MUPIROCIN			1 TUBE/30
NABUMETONE	500MG	2	180/90
NABUMETONE	750MG	2	180/90
NARATRIPTAN			12/30
NASACORT AERS	55 MCG	4 SPRAYS	9.3/25
NASONEX	50MCG	4 SPRAYS	17/30
NATROBA		120ML	1 bottle/30
NAYZILAM	All Strengths		5/30
NEUPOGEN INJ	300MCG/ML		10/30
NEUPOGEN INJ	480MCG/1.6		16/30
NEUPOGEN INJ	300MCG/.5ML		5/30
NEUPOGEN INJ	480MCG/.8ML		8/30
NEURONTIN	300MG	9	315/35
NEURONTIN	600MG	9	315/35
NEXIUM	20MG	1	35/35
NEXIUM	40MG	2	70/35
NEXIUM SUS	All Strengths	1	30/30
NIFEDIPINE CR	90MG	1	90/90
NIFEDIPINE ER	60MG	1	90/90
NIFEDIPINE ER	30MG	1	90/90
NIFEDIPINE ER	60MG	1	90/90
Drug Name	Strength	Limit/Day	Limit/Days
RELPAX	All Strengths		12/30
REMODULIN	All Strengths		1 MDV/30
RESTORIL	7.5MG		10/30
RESTORIL	15MG		10/30
RESTORIL	30MG		10/30
RETIN-A		1 TUBE	1 TUBE/30
REVLIMID	All Strengths	1	35/35
REYVOW	All Strengths		4/30
RHINOCORT AQ	32MCG	8 SPRAYS	18/30
REFRESH PLUS		15 ML	1 bottle/30
REFRESH PLUS		30 ML	2 bottles/30
REFRESH TEARS		15 ML	1 bottle/30
REFRESH TEARS		30 ML	2 bottles/30
RESCULA			2 bottles/35

ODOMZO	200mg	1	30/30
OLMESARTAN	All Strengths	1	90/90
OLANZAPINE	2.5MG	3	270/90
OLANZAPINE	5MG	3	270/90
OLANZAPINE	7.5MG	3	270/90
OLANZAPINE	10MG	3	270/90
OLANZAPINE	15MH	2	180/90
OLANZAPINE	20MG	1.5	135/90
OLANZAPINE ODT	All Strengths	1	90/90
OMEPRAZOLE	10MG	1	90/90
OMEPRAZOLE	20MG	1	90/90
OMEPRAZOLE	40MG	1	90/90
OMNARIS	50MCG	4 sprays	12.5/30
ONGLYZA	All Strengths	1	35/35
OPSUMIT	All Strengths	1	35/35
ORUVAIL	100MG	2	70/35
ORUVAIL	200MG	1	35/35
OXAPROZIN	600MG	2	180/90
OXYCODONE ER	10,20,40MG	2	70/35
OXYCODONE ER	80MG	4	140/35
OXYCONTIN**	10,20,30,40MG	2	70/35
OXYCONTIN**	80MG	4	140/35
PANTOPRAZOLE	All Strengths	1	90/90
PAROXETINE	10MG	2	180/90
PAROXETINE	20MG	2	180/90
PAXIL	10MG	1.5	53/35
PAXIL	20MG	1	35/35
PEGASYS KIT		KIT	1/28
PLAN B			2/15 or 4/30
PLENDIL	2.5MG	1	35/35
PLENDIL	5MG	1.5	53/35
PRAVACHOL	10MG	1	35/35
PRAVACHOL	20MG	1	35/35
PRAVACHOL	40MG	1	35/35
PRAVACHOL	80MG	1	35/35
PRAVASTATIN	10MG	1	35/35
PRAVASTATIN	20MG	1	35/35
PRAVASTATIN	40MG	2	180/90
PRAVASTATIN	80MG	1	35/35
PREVPAC MIS	500MG-30MG		14/30
PRILOSEC OTC	20MG	2	168/84
PRINIVIL	2.5MG	1	35/35
PRINIVIL	5MG	1	35/35
PRINIVIL	10MG	1.5	53/35
PRINIVIL	20MG	1.5	53/35
PRINZIDE	10-12.5	1	35/35
PROAIR HFA	90mcg	12 INHALATIONS	17/34
PROTONIX	20MG	2	70/35
PROTONIX	40MG	2	70/35
PROZAC	10MG	1.5	53/35
PULMICORT	200MCG	8 INHALATIONS	1/25
PULMICORT FLEX	All Strengths	8 Inhalations	2/30
QUETIAPINE	25MG	3	270/90
QUETIAPINE	50MG	3	270/90
QUETIAPINE	100MG	3	270/90
QUETIAPINE	200MG	3	270/90
QUINAPRIL	5MG	1	90/90
QUINAPRIL	10MG	1	90/90
QUINAPRIL	20MG	1	90/90
QVAR AERS	All Strengths	8 Inhalations	14.6/25
RANITIDINE SYRUP***	15MG/ML	20ML	700ML/35
RELAFEN	500MG	2	70/35
RELAFEN	750MG	2	70/35
REMERON	15MG	1.5	53/35
Drug Name	Strength	Limit/Day	Limit/Days
SULAR	10MG	1.5	53/35
SULAR	20MG	1	35/35
SUMATRIPTAN PEN INJ	All Strengths		12/30
SUMATRIPTAN NASAL SPRAY	All Strengths		12/30
SUMATRIPTAN SYRINGE	All Strengths		12/30
SUMATRIPTAN TAB	All Strengths		12/30
SYNVISC INJ	8MG/ML		2/30
SYRINGES		10	1000/100
TAFINLAR	50MG	6	210/35
TAFINLAR	75MG	4	140/35

REYATAZ	All Strengths	1	35/35
RISPERDAL	0.5MG	1.5	53/35
RISPERDAL	0.25MG	1.5	53/35
RISPERDAL	1MG	1.5	53/35
RISPERDAL	2MG	1.5	53/35
RISPERDAL	3MG	2	70/35
RISPERDAL	4MG	2	70/35
RISPERDAL INJ	25MG		2/28
RISPERDAL INJ	37.5		2/28
RISPERDAL INJ	50MG		2/28
RISPERDAL M-TAB	0.5MG	1.5	53/35
RISPERDAL M-TAB	1MG	1.5	53/35
RISPERDAL M-TAB	2MG	4	140/35
RISPERDAL SOL.	1MG/ML	8ML	280/35
RISPERIDONE	0.5MG	3	270/90
RISPERIDONE	0.25MG	3	270/90
RISPERIDONE	1MG	3	270/90
RISPERIDONE	2MG	3	270/90
RISPERIDONE	3MG	2	180/90
RISPERIDONE	4MG	2	180/90
RISPERIDONE SOL.	1MG/ML	8ML	280/35
RITALIN LA	All Strengths	1	35/35
RITALIN LA	30mg	2	70/35
SAVELLA	All Strengths	2	70/35
SEREVENT DISKUS	50MCG	2 INHALATIONS	60/30
SEROQUEL	100MG		45/30
SEROQUEL XR	150MG	1	35/35
SEROQUEL XR	200MG	1	35/35
SEROQUEL XR	300MG	2	70/35
SEROQUEL XR	400MG	2	70/35
SERTRALINE	25MG	3	270/90
SERTRALINE	50MG	3	270/90
SERTRALINE	100MG	3	270/90
SIMVASTATIN	5MG	1	35/35
SIMVASTATIN	10MG	1.5	53/35
SIMVASTATIN	20MG	1.5	53/35
SIMVASTATIN	40MG	1.5	53/35
SIMVASTATIN	80MG	1	35/35
SINGULAIR	4MG	1	35/35
SINGULAIR	5MG	1	35/35
SINGULAIR	10MG	1	35/35
SONATA	5MG		12/34
SONATA	10MG		12/34
SPIRIVA	HANDIHLR	1 INHALTION	30/30
SPORANOX SOL	10MG/ML	10ML/ML	300cc/30
SPORANOX PULSEPAK	F		30/30
SPORANOX	100MG		30/30
STADOL INJ	1MG/ML		9/35
STADOL INJ	2MG/ML		9/35
STRATTERA	All Strengths	1	35/35
SUPRAX	400MG	1	1/7

Drug Name	Strength	Limit/Day	Limit/Days
XOPENEX HFA		12 INHALATIONS	2 INHALERS/34
XOPENEX NEB		12CC	408/34
ZALEPLON	All Strengths		30/30
ZECUITY	6.5		4/28
ZEMBRACE	All Strengths		3boxes/30
ZESTORETIC	10-12.5	1	35/35
ZESTRIL	2.5MG	1	35/35
ZESTRIL	5MG	1	35/35
ZESTRIL	10MG	1.5	53/35
ZESTRIL	20MG	1.5	53/35
ZETONNA	37MCG	2	60/30
ZIPRASIDONE	20MG	3	270/90
ZIPRASIDONE	40MG	3	270/90
ZOCOR	5MG	1	35/35
ZOCOR	10MG	1.5	53/35
ZOCOR	20MG	1.5	53/35
ZOCOR	40MG	1.5	53/35
ZOFRAN*	4MG	3	90/30
ZOFRAN*	8MG	1.5	45/30
ZOFRAN*	24MG	0.5	15/30
ZOFRAN*	4MG/5ML	15ML	450/30
ZOLMITRIPTAN TAB	All Strengths		12/30

Pain Management Policy

[Back to Index](#)

Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 300 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.

However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.

The following are general exceptions: pain associated with cancer treatment, end-of-life and hospice care, palliative care, and symptoms related to HIV/AIDS. Per MaineCare criteria, the diagnosis of cancer must be written on the prescription. A palliative care exception for any MaineCare opioid prescription will require prior authorization (PA) with appropriate clinical documentation.

Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.

An MME conversion chart is available at www.mainearepd.org. Click on "General Pharmacy Info."